arXiv:1210.6931v1 [math.OA] 25 Oct 2012

DYNAMICAL SYSTEMS ASSOCIATED TO SEPARATED GRAPHS, GRAPH ALGEBRAS, AND PARADOXICAL DECOMPOSITIONS

PERE ARA AND RUY EXEL

Abstract. We attach to each finite bipartite separated graph (E,C) a partial dynamical system $(\Omega(E,C),\mathbb{F},\theta)$, where $\Omega(E,C)$ is a zero-dimensional metrizable compact space, $\mathbb{F}$ is a finitely generated free group, and θ is a continuous partial action of $\mathbb{F}$ on $\Omega(E,C)$. The full crossed product C*-algebra $\mathcal{O}(E,C)=C(\Omega(E,C))\rtimes_{\theta^*}\mathbb{F}$ is shown to be a canonical quotient of the graph C*-algebra $C^*(E,C)$ of the separated graph (E,C). Similarly, we prove that, for any $*$-field K, the algebraic crossed product $L_K^{\mathrm{ab}}(E,C)=C_K(\Omega(E,C))\rtimes_{\theta^*}^{\mathrm{alg}}\mathbb{F}$ is a canonical quotient of the Leavitt path algebra $L_K(E,C)$ of (E,C). The monoid $\mathcal{V}(L_K^{\mathrm{ab}}(E,C))$ of isomorphism classes of finitely generated projective modules over $L_K^{\mathrm{ab}}(E,C)$ is explicitly computed in terms of monoids associated to a canonical sequence of separated graphs. Using this, we are able to construct an action of a finitely generated free group $\mathbb{F}$ on a zero-dimensional metrizable compact space Z such that the type semigroup $S(Z,\mathbb{F},\mathbb{K})$ is not almost unperforated, where $\mathbb{K}$ denotes the algebra of clopen subsets of Z. Finally we obtain a characterization of the separated graphs (E,C) such that the canonical partial action of $\mathbb{F}$ on $\Omega(E,C)$ is topologically free.

Contents

1. Introduction 2
2. Preliminary definitions 6
3. Multiresolutions 8
4. Bipartite separated graphs 11
5. The main construction 14
6. Dynamical systems 23
7. The type semigroup 28
8. Descriptions of the space $\Omega(E,C)$. 36
9. Examples 43
10. Topologically free actions 50
Acknowledgements 57
References 57

Date: October 26, 2012.

2000 *Mathematics Subject Classification.* Primary 16D70, 46L35; Secondary 06A12, 06F05, 46L80.

Key words and phrases. Graph algebra, dynamical system, refinement monoid, nonstable K-theory, partial representation, partial action, crossed product, condition (L).

The first named author was partially supported by DGI MICIIN-FEDER MTM2011-28992-C02-01, and by the Comissionat per Universitats i Recerca de la Generalitat de Catalunya. The second named author was partially supported by CNPq.

1. Introduction

In their seminal paper [18], J. Cuntz and W. Krieger gave a dynamical interpretation of the Cuntz-Krieger algebras associated to square $\{0,1\}$-matrices in terms of the corresponding subshifts of finite type. Since then, there has been a long and fruitful interaction between the theory of combinatorially defined C*-algebras and dynamics, see [21], [38], [37], [25], [36], [12], [28], [11], [44], [52], [48], [47], [46], [27], [24].

In [22, Section 5], the second named author describes any Cuntz-Krieger C*-algebra as a crossed product of a commutative C*-algebra by a partial action of a non-abelian free group. The same applies to graph C*-algebras, which are generalizations of Cuntz-Krieger algebras.

A wider class of graph algebras has been introduced in [8] and [7]. These algebras are attached to *separated graphs*, where a separated graph is a pair (E,C) consisting of a directed graph E and a set $C=\bigsqcup_{v\in E^0} C_v$, where each C_v is a partition of the set of edges whose terminal vertex is v. Their associated graph C*-algebras $C^*(E,C)$ and Leavitt path algebras $L_K(E,C)$ (for an arbitrary field K), provide generalizations of the usual graph C*-algebras ([45]) and Leavitt path algebras ([2], [10]) associated to directed graphs, although these algebras behave quite differently from the usual graph algebras because the range projections associated to different edges need not commute. One motivation for their introduction was to provide graph-algebraic models for the Leavitt algebras $L_K(m,n)$ of [39], and the C*-algebras $U^{\text{nc}}_{m,n}$ studied L. Brown [15] and McClanahan [40], [41], [42]. Another motivation was to obtain graph algebras whose structure of projections is as general as possible. Thanks to deep results due to G. Bergman [16] [17], it is possible to show that the natural map $M(E,C)\to\mathcal{V}(L_K(E,C))$ from a monoid $M(E,C)$ naturally associated to (E,C) to the monoid $\mathcal{V}(L_K(E,C))$ of isomorphism classes of finitely generated projective right modules over $L_K(E,C)$ is an isomorphism, see [8, Theorem 4.3]. Since each conical abelian monoid M is isomorphic to a monoid of the form $M(E,C)$ for a suitable separated graph (E,C) ([8, Proposition 4.4]), we conclude that Leavitt path algebras of separated graphs realize all the conical abelian monoids. (By the results of [10], this is not true for the class of Leavitt path algebras of non-separated graphs.) The first author and Ken Goodearl raised in [7] the problem of whether the natural map $M(E,C)\to\mathcal{V}(C^*(E,C))$ is an isomorphism (see also [4, Section 6] for a brief discussion on this problem).

Recall that a set S of partial isometries in a $*$-algebra $\mathcal{A}$ is said to be *tame* [23, Proposition 5.4] if every element of $U=\langle S\cup S^*\rangle$, the multiplicative semigroup generated by $S\cup S^*$, is a partial isometry, and the set $\{e(u)\mid u\in U\}$ of final projections of the elements of U is a commuting set of projections of $\mathcal{A}$. A main difficulty in working with $C^*(E,C)$ and $L_K(E,C)$ is that, in general, the generating set of partial isometries of these algebras is not tame. This is not the case for the usual graph algebras, where it can be easily shown that the generating set of partial isometries is tame.

The main objective of the present paper is to associate a partial dynamical system to any finite bipartite separated graph (E,C), and to uncover some of the fundamental properties of the crossed products corresponding to it. To a large extent, this task can be made simultaneously in the purely algebraic category and in the C*-algebraic category. However, several analytic aspects, such as the study of exactness, nuclearity and reduced crossed products will need more specific tools. This dynamical system is described by the action θ of a finitely generated free non-abelian group $\mathbb{F}$, whose rank coincides with the number of edges of E, on a zero-dimensional metrizable compact space $\Omega(E,C)$. The corresponding C*-algebra full crossed product $\mathcal{O}(E,C) := C(\Omega(E,C)) \rtimes_{\theta^*} \mathbb{F}$ is shown to coincide with a canonical quotient of the graph C*-algebra $C^*(E,C)$ of the separated graph (E,C) (Corollary 6.12). Namely we obtain an isomorphism

$$C(\Omega(E,C)) \rtimes_{\theta^*} \mathbb{F} \cong C^*(E,C)/J$$

where J is the closed ideal of $C^*(E,C)$ generated by all the commutators $[e(\gamma), e(\mu)]$, where γ, μ belong to the multiplicative semigroup U of $C^*(E,C)$ generated by the canonical partial isometries, corresponding to $E^1 \cup (E^1)^*$. Note that $E^1 \cup (E^1)^*$ becomes a tame set of partial isometries in $\mathcal{O}(E,C)$. The ideal J is the closure of an increasing union of ideals J_n, with J_n the ideal generated by all the commutators $[e(\gamma), e(\mu)]$, where γ, μ can be expressed as products of $\leq n$ canonical generators. We show in Section 5 that there is a canonical sequence $\{(E_n, C^n)\}$ of finite bipartite separated graphs such that $C^*(E,C)/J_n \cong C^*(E_n, C^n)$ for all n. Consequently the C*-algebra $\mathcal{O}(E;C) = C^*(E,C)/J$ is isomorphic to an inductive limit $\varinjlim_n C^*(E_n, C^n)$ with surjective transition maps. This gives an approximation result of the dynamical C*-algebra $\mathcal{O}(E,C)$ by graph C*-algebras of separated graphs. We also consider in Section 10 a reduced C*-algebra $\mathcal{O}^r(E,C)$, which is defined in terms of the *reduced crossed product* of $C(\Omega(E,C))$ by $\mathbb{F}$. These C*-algebras have been studied in detail in [5] in the case where $(E,C) = (E(m,n), C(m,n))$ is the separated graph whose graph C*-algebra is Morita-equivalent to MacClanahan's algebras $U^{\mathrm{nc}}_{m,n}$ (see Example 9.3 for the definition), under the notations $\mathcal{O}_{m,n}$ and $\mathcal{O}^r_{m,n}$. They are attached to the so-called universal (m,n)-dynamical system, see [5, Theorem 3.8]. It is shown in [5] that the natural map $\mathcal{O}_{m,n} \to \mathcal{O}^r_{m,n}$ is not injective.

A similar result holds when we work in the category of $*$-algebras, obtaining a corresponding isomorphism

$$C_K(\Omega(E,C)) \rtimes^{\mathrm{alg}}_{\theta^*} \mathbb{F} \cong L_K(E,C)/J \cong \varinjlim L_K(E_n, C^n),$$

where $C_K(\Omega(E,C))$ is the $*$-algebra of continuous functions from $\Omega(E,C)$ to the $*$-field K endowed with the discrete topology, and the crossed product is taken in the algebraic category. We denote this quotient $*$-algebra $L_K(E,C)/J$ by $L^{\mathrm{ab}}_K(E,C)$. (Here, of course, J means the algebraic ideal generated by the commutators $[e(\gamma), e(\mu)]$, for $\gamma, \mu \in U$.) This leads to a monoid isomorphism $\mathcal{V}(L^{\mathrm{ab}}_K(E,C)) \cong \varinjlim_n \mathcal{V}(L_K(E_n, C^n)) \cong \varinjlim_n M(E_n, C^n)$. Conjecturally, the same monoid $\varinjlim_n M(E_n, C^n)$ would compute the nonstable K-theory of $\mathcal{O}(E,C)$. The monoid $\mathfrak{R}(E,C) := \varinjlim M(E_n, C^n)$ is a refinement monoid (see Section 2 for the definition) and the natural map $M(E,C) \to \mathfrak{R}(E,C)$ gives a canonical refinement of $M(E,C)$ (Lemma 4.5). This condition implies in particular that the map $M(E,C) \to \mathfrak{R}(E,C)$ is an order-embedding.

We remark that there is no loss of generality in assuming that our separated graphs are bipartite, as we show in Proposition 9.1 that every graph algebra of a separated graph is Morita-equivalent to a graph algebra of a bipartite separated graph.

As a biproduct of our results, we are able to answer a question raised in recent papers by Kerr [34], Kerr and Nowak [35], and Rørdam and Sierakowski [50]. The statement of this question is unrelated to graph algebras but, using our new theory, we can give a complete answer to it. The classical Tarski's Theorem ([54, Corollary 9.2]) asserts that, for an action of a group G on a set X, a subset E of X is not G-paradoxical if and only if there is a finitely additive G-invariant measure $\mu\colon \mathcal{P}(X) \to [0,\infty]$ such that $\mu(E) = 1$. This result follows from special properties of the *type semigroup* $S(X,G)$ of the action. Observe that the set of finitely additive G-invariant measures μ as above is precisely the state space of $(S(X,G),[E])$.

If G is a discrete group acting by continuous transformations on a topological space X and $\mathbb{D}$ is a G-invariant subalgebra of subsets of X, one may restrict the equidecomposability relation to subsets in $\mathbb{D}$, obtaining a "relative" type semigroup $S(X,G,\mathbb{D})$, as in [34], [35], [50]. Of particular interest is the case where X is a zero-dimensional metrizable compact space and $\mathbb{D}$ is the subalgebra $\mathbb{K}$ of clopen subsets of X. It has been asked in the above-mentioned papers (cf. [50, page 285], [34, Question 3.10]) whether the analogue of Tarski's Theorem holds in this more general context, that is, whether for $E \in \mathbb{K}$ we have that $2[E] \nleq [E]$ in $S(X,G,\mathbb{K})$ if and only if the state space of $(S(X,G,\mathbb{K}),[E])$ is non-empty. We show in Corollary 7.12 that the answer to this question is negative. Indeed, our Theorem 7.11 implies that $S(X,G,\mathbb{K})$ does not satisfy in general any cancellation or order-cancellation law, since, given any finitely generated conical abelian monoid M we can find a triple $(X,G,\mathbb{K})$, with X a 0-dimensional metrizable compact space, such that there is an order-embedding of M into $S(X,G,\mathbb{K})$. Our methods give in a natural way *partial actions* of discrete groups on totally disconnected spaces. We use globalization results from [1] to reach the desired goal with *global actions*.

We also obtain a characterization of the finite bipartite separated graphs (E,C) such that the canonical action of the free group $\mathbb{F}$ on the space $\Omega(E,C)$ is topologically free. It turns out that this is equivalent to a version, adapted to separated graphs, of the well-known condition (L) for graphs, which is the condition that every cycle has an entry ([37], [25, Definition 12.1]), see Theorem 10.5. Using this result and known facts on crossed products by partial actions from [26] we show that, if (E,C) satisfies condition (L), then the *reduced crossed product* C*-algebra $\mathcal{O}^r(E,C) = C(\Omega(E,C)) \rtimes^r \mathbb{F}$ satisfies condition (SP), that is, every nonzero hereditary subalgebra contains a nonzero projection, see Theorem 10.9. A corresponding result is also shown for the algebra $L_K^{\mathrm{ab}}(E,C)$ (Theorem 10.10 and Remark 10.11).

Many properties of these new classes of algebras remain to be investigated. We mention the computation of K-theoretic invariants in terms of the separated graph (E,C), the study of the structure of ideals, and the study of exactness and nuclearity. Topologically free minimal orbits of the topological space $\Omega(E,C)$ with respect to the canonical action of the free group will surely provide examples of simple C*-algebras with exotic properties. In a different direction, the algebras constructed in the present paper represent a further step in the strategy initiated in [8] to construct von Neumann regular rings and exchange rings with prescribed $\mathcal{V}$-monoids (see [3] for a survey on this problem). This will be discussed in more detail in [9], where the refinement monoid $\mathfrak{R}(E,C) = \varinjlim M(E_n, C^n)$ associated to the separated graph (E,C) described in Example 9.6 will be realized as the $\mathcal{V}$-monoid of an exchange K-algebra obtained as a universal localization of the corresponding algebra $L_K^{\mathrm{ab}}(E,C)$. The latter algebra is Morita equivalent to the algebra of the *free monogenic inverse semigroup* (see Example 9.6).

Contents. We now explain in more detail the contents of this paper. In Section 2 we recall the basic definitions needed for our work, coming from the papers [7] and [8]. Section 3 contains some preparatory material concerning graph theory and semigroup theory. In particular we introduce the crucial concept of a multiresolution of a separated graph (E,C) at a set of vertices V of E. This is a modification of the concept of resolution of a separated graph at a set of vertices, which has been introduced in [8, Section 8]. As in [8], the aim to introduce this concept is to give embeddings of the monoid $M(E,C)$ associated to a separated graph (E,C) into a refinement monoid corresponding to another separated graph (F,D), given in general as the limit of an infinite sequence of separated graphs obtained by successive applications of the resolution process. The multiresolution process differs from the resolution process in that it involves at once all the sets in the corresponding partitions C_v, $v \in V$, and so it does not depend on the particular choice of pairs $X, Y \in C_v$, for $v \in V$. In Section 4, we define, for a given finite bipartite separated graph (E,C), a canonical sequence of finite separated graphs (F_n, D^n), obtained by successive applications of the multiresolution process, in such a way that the union $(F_\infty, D^\infty) = \bigcup_{n=0}^\infty (F_n, D^n)$ satisfies that $M(F_\infty, D^\infty)$ is a refinement of $M(E,C)$. The canonical sequence $\{(E_n, C^n)\}$ of finite bipartite separated graphs obtained as the *sections* of (F_∞, D^∞) is of special importance. On one hand, it is shown in Lemma 4.5 that there are natural connecting homomorphisms $\iota_n \colon M(E_n, C^n) \to M(E_{n+1}, C^{n+1})$ such that $M(F_\infty, D^\infty) \cong \varinjlim_n (M(E_n, C^n), \iota_n)$, so that the map $M(E,C) \to \varinjlim_n (M(E_n, C^n), \iota_n)$ is a refinement of $M(E,C)$. On the other hand, it is shown in Theorem 5.6 that there is a natural isomorphism $L_K(E,C)/J_n \cong L_K(E_n, C^n)$, where J_n is the ideal of $L_K(E,C)$ generated by all the commutators $[e(\gamma), e(\mu)]$, where γ, μ are products of $\leq n$ canonical generators from $E^1 \cup (E^1)^*$. A similar result holds for the corresponding graph C*-algebras. The notations $L^{\mathrm{ab}}(E,C)$ and $\mathcal{O}(E,C)$ are introduced in 5.7. The topological space $\Omega(E,C)$ is also introduced in Section 5 as the spectrum of the canonical commutative AF-subalgebra $\mathcal{B}_0$ of $\mathcal{O}(E,C)$.

We develop in Section 6 the crossed product structure of the algebras $L_K^{\mathrm{ab}}(E,C)$ and $\mathcal{O}(E,C)$. This is done in a soft way, using the universal properties of the involved objects. The topological space $\Omega(E,C)$ is shown in Corollary 6.11 to be the *universal* (E,C)-*dynamical system.* This gives a different proof of [5, Theorem 3.8], where the particular case of (m,n)-dynamical systems is considered. Moreover the natural partial action of the free group $\mathbb{F} = \mathbb{F}\langle E^1 \rangle$ on $\Omega(E,C)$ induces $*$-isomorphisms $L_K^{\mathrm{ab}}(E,C) \cong C_K(\Omega(E,C)) \rtimes_\alpha \mathbb{F}$ and $\mathcal{O}(E,C) \cong C(\Omega(E,C)) \rtimes_\alpha \mathbb{F}$ (Corollary 6.12). Section 7 is devoted to the study of the type semigroups. Our main result is Theorem 7.11, where it is shown that, given any finitely generated abelian conical monoid M there exists a 0-dimensional metrizable compact space Z and an action of a finitely generated free group $\mathbb{F}$ on it so that M order-embeds in the type semigroup $S(Z, \mathbb{F}, \mathbb{K})$. This uses the machinery from Sections 3–6 plus Abadie's globalization results [1]. In Section 8 we develop descriptions of the space $\Omega(E, C)$ and the canonical partial action of the free group $\mathbb{F}$ on it. There are two different descriptions, one in terms of configurations and another in terms of certain "choice functions", called here E-functions. We show in Theorem 8.3 that the basic clopen sets corresponding to the vertices in the odd layers of the graph F_∞ (introduced in Construction 4.4) correspond precisely to the partial E-functions. Several examples are presented in Section 9. Apart from the motivational example leading to the (m, n)-dynamical system (Example 9.3), we consider various interesting examples which are connected with constructions already investigated in disparate contexts. So for example we recover Truss example from [53] of a G-space X with failure of the 2-cancellation law $2x = 2y \implies x = y$ in the type semigroup (Example 9.5), a separated graph (E, C) such that the algebra $\mathcal{O}(E, C)$ is Morita-equivalent to the C*-algebra of the free monogenic inverse semigroup [33] (Example 9.6), and another separated graph (F, D) such that $\mathcal{O}(F, D)$ (respectively $L_K^{\mathrm{ab}}(F, D)$) is Morita-equivalent to the group C*-algebra of the lamplighter group $C^*(\mathbb{Z}_2 \wr \mathbb{Z})$ (respectively the group algebra $K[\mathbb{Z}_2 \wr \mathbb{Z}]$). Finally, Section 10 contains our characterization of the finite bipartite separated graphs (E, C) such that the canonical action of the free group group $\mathbb{F}$ on the space $\Omega(E, C)$ is topologically free and the consequences for the algebraic structure of $\mathcal{O}(E, C)$ and $L_K^{\mathrm{ab}}(E, C)$.

2. Preliminary definitions

The concept of separated graph, introduced in [8], plays a vital role in our construction. In this section, we will recall this concept and we will also recall the definitions of the *monoid associated to a separated graph*, the *Leavitt path algebra* and the *graph C*-algebra* of a separated graph.

We will use the reverse notation as in [8] and [7], but in agreement with the one used in [5], and in the book [45].

Definition 2.1. ([8]) A *separated graph* is a pair (E, C) where E is a graph, $C = \bigsqcup_{v \in E^0} C_v$, and C_v is a partition of $r^{-1}(v)$ (into pairwise disjoint nonempty subsets) for every vertex v. (In case v is a source, we take C_v to be the empty family of subsets of $r^{-1}(v)$.)

If all the sets in C are finite, we say that (E, C) is a *finitely separated* graph. This necessarily holds if E is column-finite (that is, if $r^{-1}(v)$ is a finite set for every $v \in E^0$.)

The set C is a *trivial separation* of E in case $C_v = \{r^{-1}(v)\}$ for each $v \in E^0 \setminus \mathrm{Source}(E)$. In that case, (E, C) is called a *trivially separated graph* or a *non-separated graph*.

The following definition gives the Leavitt path algebra $L_K(E, C)$ as a universal object in the category of $*$-algebras over a fixed field with involution K. The underlying algebra is naturally isomorphic to the algebra defined in [8, Definition 2.2] in case every vertex is either the range vertex or the source vertex of some edge in E.

Definition 2.2. Let $(K,*)$ be a field with involution. The *Leavitt path algebra of the separated graph* (E,C) with coefficients in the field K, is the $*$-algebra $L_K(E,C)$ with generators $\{v,e \mid v\in E^0, e\in E^1\}$, subject to the following relations:

(V) $vv'=\delta_{v,v'}v$ and $v=v^*$ for all $v,v'\in E^0$,

(E) $r(e)e=es(e)=e$ for all $e\in E^1$,

(SCK1) $e^*e'=\delta_{e,e'}s(e)$ for all $e,e'\in X$, $X\in C$, and

(SCK2) $v=\sum_{e\in X}ee^*$ for every finite set $X\in C_v$, $v\in E^0$.

The Leavitt path algebra $L_K(E)$ is just $L_K(E,C)$ where $C_v=\{r^{-1}(v)\}$ if $r^{-1}(v)\neq\emptyset$ and $C_v=\emptyset$ if $r^{-1}(v)=\emptyset$. An arbitrary field can be considered as a field with involution by taking the identity as the involution. However, our "default" involution over the complex numbers $\mathbb{C}$ will be the complex conjugation.

We now recall the definition of the graph C*-algebra $C^*(E,C)$, introduced in [7].

Definition 2.3. The *graph C*-algebra* of a separated graph (E,C) is the C*-algebra $C^*(E,C)$ with generators $\{v,e \mid v\in E^0,\ e\in E^1\}$, subject to the relations (V), (E), (SCK1), (SCK2). In other words, $C^*(E,C)$ is the enveloping C*-algebra of $L(E,C)$.

In case (E,C) is trivially separated, $C^*(E,C)$ is just the classical graph C*-algebra $C^*(E)$. There is a unique *-homomorphism $L_{\mathbb{C}}(E,C)\to C^*(E,C)$ sending the generators of $L(E,C)$ to their canonical images in $C^*(E,C)$. This map is injective by [7, Theorem 3.8(1)].

If no confusion can arise, we will suppress the field K from our notation.

Since both $L_K(E,C)$ and $C^*(E,C)$ are universal objects with respect to the same sets of generators and relations, many of the constructions of this paper apply in the same way to either class. A remarkable difference is that we *know* exactly what is the structure of the monoid $\mathcal{V}(L_K(E,C))$ for any separated graph (E,C), but we still do not know the structure of the monoid $\mathcal{V}(C^*(E,C))$, although it is conjectured in [7] that the natural map $L_{\mathbb{C}}(E,C)\to C^*(E,C)$ induces an isomorphism $\mathcal{V}(L_{\mathbb{C}}(E,C))\to\mathcal{V}(C^*(E,C))$. See [4, Section 6] for a short discussion on this problem.

Recall that for a unital ring R, the monoid $\mathcal{V}(R)$ is usually defined as the set of isomorphism classes $[P]$ of finitely generated projective (left, say) R-modules P, with an addition operation given by $[P]+[Q]=[P\oplus Q]$. For a nonunital version, see Definition [8, Definition 10.8].

For arbitrary rings, $\mathcal{V}(R)$ can also be described in terms of equivalence classes of idempotents from the ring $M_\infty(R)$ of $\omega\times\omega$ matrices over R with finitely many nonzero entries. The equivalence relation is *Murray-von Neumann equivalence*: idempotents $e,f\in M_\infty(R)$ satisfy $e\sim f$ if and only if there exist $x,y\in M_\infty(R)$ such that $xy=e$ and $yx=f$. Write $[e]$ for the equivalence class of e; then $\mathcal{V}(R)$ can be identified with the set of these classes. Addition in $\mathcal{V}(R)$ is given by the rule $[e]+[f]=[e\oplus f]$, where $e\oplus f$ denotes the block diagonal matrix $\left(\begin{smallmatrix} e & 0\\ 0 & f\end{smallmatrix}\right)$. With this operation, $\mathcal{V}(R)$ is an abelian monoid, and it is *conical*, meaning that $a+b=0$ in $\mathcal{V}(R)$ only when $a=b=0$. Whenever A is a C^*-algebra, the monoid $\mathcal{V}(A)$ agrees with the monoid of equivalence classes of projections in $M_\infty(A)$ with respect to the equivalence relation given by $e\sim f$ if and only if there is a partial isometry w in $M_\infty(A)$ such that $e=ww^*$ and $f=w^*w$; see [13, 4.6.2 and 4.6.4] or [49, Exercise 3.11].

We will need the definition of $M(E,C)$ only for finitely separated graphs. The reader can consult [8] for the definition in the general case. Let (E,C) be a finitely separated graph, and let $M(E,C)$ be the abelian monoid given by generators a_v, $v \in E^0$, and relations $a_v = \sum_{e \in X} a_{s(e)}$, for $X \in C_v$, $v \in E^0$. Then there is a canonical monoid homomorphism $M(E,C) \to \mathcal{V}(L_K(E,C))$, which is shown to be an isomorphism in [8, Theorem 4.3]. The map $\mathcal{V}(L_{\mathbb{C}}(E,C)) \to \mathcal{V}(C^*(E,C))$ induced by the natural $*$-homomorphism $L_K(E,C) \to C^*(E,C)$ is conjectured to be an isomorphism for all finitely separated graphs (E,C) (see [7] and [4, Section 6]).

3. Multiresolutions

In this section, we will introduce the concept of mutiresolution of a finitely separated graph (E,C) at a vertex $v \in E^0$, which is closely related to the notion of resolution, studied in [8]. These computations will be heavily used in the forthcoming sections.

An (abelian) monoid M is said to be a *refinement monoid* if whenever $a+b=c+d$ in M, there exist x,y,z,t in M such that $a=x+y$ and $b=z+t$ while $c=x+z$ and $d=y+t$.

We recall the following definitions from [8]. For $X \in C$, we write $\mathbf{s}(X) := \sum_{e \in X} s(e)$.

Definition 3.1. Let F be the free abelian monoid on E^0. For $\alpha, \beta \in F$, write $\alpha \rightsquigarrow_1 \beta$ to denote the following situation:

> $\alpha = \sum_{i=1}^k u_i + \sum_{i=k+1}^m u_i$ and $\beta = \sum_{i=1}^k \mathbf{s}(X_i) + \sum_{i=k+1}^m u_i$ for some $u_i \in E^0$ and some $X_i \in C_{u_i}$, where $u_1, \dots, u_k \in E^0 \setminus \mathrm{Source}(E)$.

We view $\alpha = 0$ as an empty sum of the above form (i.e., $k=m=0$), so that $0 \rightsquigarrow_1 0$. Note also, taking $k=0$, that $\alpha \rightsquigarrow_1 \alpha$ for all $\alpha \in F$.

Definition 3.2. Assumption $(*)$: Let $v \in E^0$. Then (E,C) satisfies $(*)$ at v if for all $X,Y \in C_v$, there exists $\gamma \in F$ such that $\mathbf{s}(X) \rightsquigarrow_1 \gamma$ and $\mathbf{s}(Y) \rightsquigarrow_1 \gamma$.

This assumption only needs to be imposed when X and Y are disjoint, since otherwise $X=Y$ and the conclusion is trivially satisfied.

The separated graph (E,C) satisfies Assumption $(*)$ in case it satisfies $(*)$ at every vertex $v \in E^0$.

Assumption (*) guarantees the refinement property in the monoid $M(E,C)$:

Proposition 3.3. [8, Proposition 5.9] *Let (E,C) be a finitely separated graph. If Assumption $(*)$ holds, then the monoid $M(E,C)$ is a refinement monoid.*

In fact [8] contains a version of the above result which holds for arbitrary separated graphs (E,C).

The multiresolution of a separated graph (E,C) at a vertex v (with $|r^{-1}(v)| < \infty$) is the simultaneous resolution (by refinement) of *all* the subsets $X \in C_v$ at once. We describe this process here. This should be compared with the resolution process of [8], where only *pairs* $X, Y \in C_v$ are refined.

Definition 3.4. Let $C_v = \{X_1, \dots, X_k\}$ with each X_i a finite subset of $r^{-1}(v)$. Put $M = \prod_{i=1}^{k} |X_i|$. Then the multiresolution of (E, C) at v is the separated graph (E_v, C^v) with

$$E_v^0 = E^0 \sqcup \{v(x_1, \dots, x_k) \mid x_i \in X_i, i = 1, \dots, k\},$$

and with $E_v^1 = E^1 \sqcup \Lambda$, where Λ is a new set of arrows defined as follows. For each $x_i \in X_i$, we put $M/|X_i|$ new arrows $\alpha^{x_i}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_k)$, $x_j \in X_j$, $j \neq i$, with

$$r(\alpha^{x_i}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_k)) = s(x_i), \text{ and } s(\alpha^{x_i}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_k)) = v(x_1, \dots, x_k).$$

For a vertex $w \in E^0$, define the new groups at w as follows. These groups are indexed by the edges $x_i \in X_i$, $i = 1, \dots, k$, such that $s(x_i) = w$. For one such x_i, set

$$X(x_i) = \{\alpha^{x_i}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_k) \mid x_j \in X_j, j \neq i\}.$$

Then

$$(C^v)_w = C_w \sqcup \{X(x_i) \mid x_i \in X_i, s(x_i) = w, i = 1, \dots, k\}.$$

The new vertices $v(x_1, \dots, x_k)$ are sources in E_v.

Definition 3.5. Let $V \subseteq E^0$ be a set of vertices such that, for each $u \in V$, $C_u = \{X_1^u, \dots, X_{k_u}^u\}$, with each X_i^u a finite subset of $r^{-1}(u)$. Then the *multiresolution of E at V* is the separated graph (E_V, C^V) obtained by applying the above process to all vertices u in V.

Hence

$$E_V^0 = E^0 \sqcup \Big(\bigsqcup_{u \in V} \{v(x_1^u, \dots, x_{k_u}^u) \mid x_i^u \in X_i^u, i = 1, \dots, k_u\} \Big),$$

and $E_V^1 = E^1 \sqcup \Big(\bigsqcup_{u \in V} \Lambda_u \Big)$, where Λ_u is the corresponding set of arrows $\alpha^{x_i^u}(x_1^u, \dots, x_{i-1}^u, x_{i+1}^u, \dots, x_{k_u}^u)$ defined as in Definition 3.4, for each $u \in V$. The sets $(C^V)_w$, for $w \in E_V^0$, are defined just as in Definition 3.4:

$$(C^V)_w = C_w \sqcup \{X(x_i^u) \mid x_i^u \in X_i^u, s(x_i^u) = w, i = 1, \dots, k_u, u \in V\}.$$

The new vertices $v(x_1^u, \dots, x_{k_u}^u)$ are sources in E_V.

Lemma 3.6. *Let $V \subseteq E^0$ be a set of vertices such that, for each $u \in V$, $C_u = \{X_1^u, \dots, X_{k_u}^u\}$, with each X_i^u a finite subset of $r^{-1}(u)$. Then the separated graph (E_V, C^V) satisfies (*) at u for all $u \in V$.*

Proof. Observe that, for $x_i^u \in X_i^u$, we have $s(x_i^u) \to_1 \sum_{x_j^u \in X_j^u, j \neq i} v(x_1^u, \dots, x_{i-1}^u, x_i^u, x_{i+1}^u, \dots, x_{k_u}^u)$, so that

$$\mathbf{s}(X_i^u) \rightsquigarrow_1 \sum_{(x_1^u, \dots, x_{k_u}^u) \in X_1^u \times \cdots \times X_{k_u}^u} v(x_1^u, \dots, x_{k_u}^u) =: \gamma.$$

This clearly shows the result, indeed we have $\mathbf{s}(X_i^u) \rightsquigarrow_1 \gamma$, for all $i = 1, \dots, k_u$. $\square$

Let us recall the definition of unitary embedding:

Definition 3.7. Following [55], a monoid homomorphism $\psi : M \to F$ is *unitary* provided

1. ψ is injective;
2. $\psi(M)$ is *cofinal* in F, that is, for each $u \in F$ there is some $v \in M$ with $u \leq \psi(v)$;

(3) whenever $u, u' \in M$ and $v \in F$ with $\psi(u) + v = \psi(u')$, we have $v \in \psi(M)$.

Lemma 3.8. *Let (E, C) be a separated graph and let $V \subseteq E^0$ be a set of vertices such that $|r^{-1}(u)| < \infty$ for all $u \in V$. Let $\iota : (E, C) \to (E_V, C^V)$ denote the inclusion morphism, where (E_V, C^V) is the multiresolution of (E, C) at V. Then $M(\iota) : M(E, C) \to M(E_V, C^V)$ is a unitary embedding.*

Proof. The proof follows the steps of the proof of [8, Lemma 8.6].

We will sketch some of the details.

Set $\mu = M(\iota)$. For $u \in V$, set $C_u = \{X_1^u, \dots, X_{k_u}^u\}$. Let F be the free monoid on generators $a(x_1^u, \dots, x_{k_u}^u)$, for $x_i^u \in X_i^u$, $i = 1, \dots, k_u$, $u \in V$. Let M be the monoid given by generators $\{b(x_i^u) \mid x_i^u \in X_i^u, i = 1, \dots, k_u, u \in V\}$ with the relations $\sum_{x_i^u \in X_i^u} b(x_i^u) = \sum_{x_j^u \in X_j^u} b(x_j^u)$ for all $1 \le i < j \le k_u$, for all $u \in V$. There is a natural homomorphism $\psi : M \to F$ sending $b(x_i^u)$ to

$$\sum_{j \neq i} \sum_{x_j^u \in X_j^u} a(x_1^u, \dots, x_{i-1}^u, x_i^u, x_{i+1}^u, \dots, x_k^u)$$

for $x_i^u \in X_i^u$, $i = 1 \dots, k_u$, $u \in V$. Arguments similar to the ones used in the proof of [8, Lemma 8.2] give that ψ is a unitary embedding.

There is a unique homomorphism $\eta : M \to M(E, C)$ sending $b(x_i^u)$ to $[s(x_i^u)]$ for $x_i^u \in X_i^u$, $1 \le i \le k_u$, $u \in V$, and there is a unique homomorphism $\eta' : F \to M(E_V, C^V)$ sending $a(x_1^u, \dots, x_{k_u}^u) \mapsto [v(x_1^u, \dots, x_{k_u}^u)]$ for $x_i^u \in X_i^u$, $i = 1, \dots, k_u$, $u \in V$. There is a commutative diagram as follows:

$$\begin{tikzcd} M \arrow[r, "\psi"] \arrow[d, "\eta"'] & F \arrow[d, "\eta'"] \\ M(E, C) \arrow[r, "\mu"] & M(E_V, C^V) \end{tikzcd} \tag{3.1}$$

An easy adaptation of the proof of [8, Lemma 8.6] gives that (3.1) is a pushout in the category of abelian monoids. It follows from [55, Lemma 1.6] that μ is a unitary embedding, completing the proof. □

Remark 3.9. (Universal property) Let (E, C) be a separated graph and let $V \subseteq E^0$ be a set of vertices such that $|r^{-1}(u)| < \infty$ for all $u \in V$.

The pushout property appearing in the proof of the above lemma is equivalent to the following universal property of $M(E_V, C^V)$: Given a monoid homomorphism $\Phi \colon M(E, C) \to N$ and given a multiresolution $\{c(x_1^u, \dots, x_{k_u}^u) \mid x_i^u \in X_i^u\}$, $u \in V$, in N of the set of equations

$$\Phi(\mathbf{s}(X_1^u)) = \Phi(\mathbf{s}(X_2^u)) = \cdots = \Phi(\mathbf{s}(X_{k_u}^u)), \quad (u \in V)$$

(so that $\Phi(s(x_i^u)) = \sum_{j \neq i, x_j^u \in X_j^u} c(x_1^u, \dots, x_{k_u}^u)$ for all $x_i^u \in X_i^u$, $u \in V$), there exists a unique monoid homomorphism $\tilde{\Phi} \colon M(E_V, C^V) \to N$ such that $\tilde{\Phi}(v) = \Phi(v)$ for all $v \in E^0$ and $\tilde{\Phi}(v(x_1^u, \dots, x_{k_u}^u)) = c(x_1^u, \dots, x_{k_u}^u)$ for all $x_i^u \in X_i^u$, $i = 1, \dots, k_u$, $u \in V$.

Note that the above multiresolutions always exist if N is a refinement monoid.

4. Bipartite separated graphs

Definition 4.1. Let E be a directed graph. We say that E is a *bipartite directed graph* if $E^0 = E^{0,0} \sqcup E^{0,1}$, with all arrows in E^1 going from a vertex in $E^{0,1}$ to a vertex in $E^{0,0}$. To avoid trivial cases, we will always assume that $r^{-1}(v) \neq \emptyset$ for all $v \in E^{0,0}$ and $s^{-1}(v) \neq \emptyset$ for all $v \in E^{0,1}$.

A *bipartite separated graph* is a separated graph (E, C) such that the underlying directed graph E is a bipartite directed graph.

Definition 4.2. Given a finitely presented abelian conical monoid

$$M = \langle \mathcal{X} \mid \mathcal{R} \rangle ,$$

where $\mathcal{X}$ is a finite set of generators $a_1, \dots, a_n$ and $\mathcal{R}$ is a finite set of relations $\mathbf{r}_1, \dots, \mathbf{r}_m$ of the form

$$\mathbf{r}_j : \qquad \sum_{i=1}^{n} r_{ji} a_i = \sum_{i=1}^{n} s_{ji} a_i, \tag{4.1}$$

where r_{ji} and s_{ji} are non-negative integers satisfying $\sum_{i=1}^n r_{ji} > 0$ and $\sum_{i=1}^n s_{ji} > 0$ for all j, and $\sum_{j=1}^m (r_{ji} + s_{ji}) > 0$ for all i, we may associate to it a finite bipartite separated graph (E, C) such that $s^{-1}(v) \neq \emptyset$ for all $v \in E^{0,0}$ and $r^{-1}(v) \neq \emptyset$ for all $v \in E^{0,1}$, as follows (cf. [8, Proposition 4.4]:)

$$E^0 = E^{0,0} \sqcup E^{0,1}, \quad \text{with} \quad E^{0,0} = \mathcal{R}, \quad E^{0,1} = \mathcal{X}.$$

For $\mathbf{r}_j \in \mathcal{R}$ given by (4.1), we set $C_{\mathbf{r}_j} = \{X_j, Y_j\}$, where X_j has exactly r_{ji} arrows from a_i to $\mathbf{r}_j$, for $i = 1, \dots, n$, and Y_j has exactly s_{ji} arrows from a_i to $\mathbf{r}_j$, for $i = 1, \dots, n$, so that $r^{-1}(\mathbf{r}_j) = X_j \sqcup Y_j$. Then $E^1 = \bigsqcup_{j=1}^m r^{-1}(\mathbf{r}_j)$. We have an isomorphism $M(E, C) \cong M$ ([8, Proposition 4.4]).

Note that there is a bijective correspondence between the finite bipartite separated graphs (E, C) satisfying the conditions in Definition 4.1 such that $|C_v| = 2$ for all $v \in E^{0,0}$ and the finite presentations of abelian conical monoids as above. Also, recall that every finitely generated abelian semigroup is finitely presented, by Redei's Theorem (see [31] for a very simple proof), so that we can apply the above process to any finitely generated abelian conical monoid. (Of course, the finite bipartite separated graph will depend on the finite presentation, not just on the monoid.)

It is useful to introduce the following terminology:

Definition 4.3. Let M be an abelian conical monoid. A *refinement* of M is an abelian conical monoid N, together with a monoid homomorphism $\iota \colon M \to N$ such that:

(a) ι is a unitary embedding.
(b) N is a refinement monoid.
(c) For each refinement monoid P and each monoid homomorphism $\psi \colon M \to P$ there is a monoid homomorphism $\widetilde{\psi} \colon N \to P$ such that $\psi = \widetilde{\psi} \circ \iota$.

Construction 4.4. (a) Let (E,C) be a finite bipartite separated graph. We define a nested sequence of finite separated graphs (F_n,D^n) as follows. Set $(F_0,D^0)=(E,C)$. Assume that a nested sequence

$$(F_0,D^0)\subset(F_1,D^1)\subset\cdots\subset(F_n,D^n)$$

has been constructed in such a way that for $i=1,\dots,n$, we have $F_i^0=\bigsqcup_{j=0}^{i+1}F^{0,j}$ for some finite sets $F^{0,j}$ and $F_i^1=\bigsqcup_{j=0}^{i}F^{1,,j}$, with $s(F^{1,j})=F^{0,j+1}$ and $r(F^{1,j})=F^{0,j}$ for $j=1,\dots,i$. We can think of (F_n,D^n) as a union of n bipartite separated graphs. Assume that condition (*) for (F_n,D^n) holds at all vertices in $\bigsqcup_{j=0}^{n-1}F^{0,j}$. Set $V_n=F^{0,n}$, and let (F_{n+1},D^{n+1}) be the multiresolution of (F_n,D^n) at V_n. Then $F_{n+1}^0=F_n^0\bigsqcup F^{0,n+2}=\bigsqcup_{j=1}^{n+2}F^{0,j}$ and $F_{n+1}^1=F_n^1\bigsqcup F_{n+1}^{1,n+1}=\bigsqcup_{j=0}^{n+1}F^{1,j}$, with $s(F^{1,n+1})=F^{0,n+2}$ and $r(F^{1,n+1})=s(F^{1,n})=F^{0,n+1}$. Moreover, by Lemma 3.6, the separated graph (F_{n+1},D^{n+1}) satisfies condition (*) at all the vertices in $\bigsqcup_{j=0}^{n}F^{0,j}$.

(b) Let

$$(F_\infty,D^\infty)=\bigcup_{n=0}^{\infty}(F_n,D^n)\,.$$

Observe that (F_∞,D^∞) is the direct limit of the sequence $\{(F_n,D^n)\}$ in the category **FSGr** defined in [8, Definition 8.4]. Since the functor $M\colon \mathbf{FSGr}\to\mathbf{Mon}$ is continuous (see [8, 8.4, 4.1]), it follows that $M(F_\infty,D^\infty)=\varinjlim M(F_n,D^n)$. Since (F_∞,D^∞) clearly satisfies condition (*) at all its vertices, it follows from [8, Proposition 5.9] that $M(F_\infty,D^\infty)$ is a refinement monoid. Moreover , since all maps $M(F_n,D^n)\to M(F_{n+1},D^{n+1})$ are unitary embeddings by Lemma 3.6, it follows that the map $M(E,C)\to M(F_\infty,D^\infty)$ is a unitary embedding. Finally, condition (c) in Definition 4.3 follows from Remark 3.9. Hence, we have that $M(E,C)\to M(F_\infty,D^\infty)$ is a refinement of $M(E,C)$. We call (F_∞,D^∞) the *complete multiresolution* of (E,C).

(c) We define a canonical sequence (E_n,C^n) of finite bipartite separated graphs as follows:

1. Set $(E_0,C^0)=(E,C)$.
2. $E_n^{0,0}=F^{0,n}$, $E_n^{0,1}=F^{0,n+1}$, and $E_n^1=F^{1,n}$. Moreover $C_v^n=D_v^n$ for all $v\in E_n^{0,0}$ and $C_v^n=\emptyset$ for all $v\in E_n^{0,1}$.

We call the sequence $\{(E_n,C^n)\}_{n\geq 0}$ the *canonical sequence of bipartite separated graphs* associated to (E,C).

Lemma 4.5. *Let (E,C) be a finite bipartite separated graph, let (E_n,C^n) be the canonical sequence of bipartite separated graphs associated to (E,C), and let $F_\infty,D^\infty)$ be the complete multiresolution of (E,C). Then the following properties hold:*

(a) *For each $n\geq 0$, there is a natural isomorphism*

$$\varphi_n\colon M(E_{n+1},C^{n+1})\longrightarrow M((E_n)_{V_n},(C^n)^{V_n}),$$

where $V_n=E_n^{0,0}=F^{0,n}$.

(b) *For each $n \geq 0$, there is a canonical unitary embedding*

$$\iota_n \colon M(E_n, C^n) \to M(E_{n+1}, C^{n+1}).$$

(c) *The canonical inclusion $j_n \colon (E_n, C^n) \to (F_n, D^n)$ induces an isomorphism*

$$M(j_n) \colon M(E_n, C^n) \to M(F_n, D^n).$$

(d) *We have $M(F_\infty, D^\infty) \cong \varinjlim (M(E_n, C^n), \iota_n)$. Consequently, the natural map $M(E, C) \to \varinjlim (M(E_n, C^n), \iota_n)$ is a refinement of $M(E, C)$.*

Proof. To simplify the notation, write $E_{V_n} := (E_n)_{V_n}$ and $C^{V_n} = (C^n)^{V_n}$.

(a) Define $\varphi_n \colon M(E_{n+1}, C^{n+1}) \longrightarrow M(E_{V_n}, C^{V_n})$ by $\varphi_n(a_v) = a_v$ for $v \in E^0_{n+1}$. Conversely, define $\psi_n \colon M(E_{V_n}, C^{V_n}) \to M(E_{n+1}, C^{n+1})$ by $\psi_n(a_v) = a_v$ for $v \in E^0_{n+1} \subset E^0_{V_n}$ and $\psi_n(a_v) = \mathbf{s}(X)$ for $X \in C_v$ if $v \in E^{0,0}_n$. We have to show that ψ_n is a well-defined monoid homomorphism. There are no relations at the vertices at $E^{0,1}_{n+1}$, both in $M(E_{n+1}, C^{n+1})$ and in $M(E_{V_n}, C^{V_n})$, because these vertices are sources in both graphs. It is clear that the relations at vertices v in $E^{0,0}_{n+1}$ are preserved by ψ_n, because $C^{n+1}_v = C^{V_n}_v$ for these vertices. For a vertex $v \in E^{0,0}_n$, take $X, Y \in C_v$. Setting $C_v = \{X_1, \dots, X_s\}$, there are $1 \leq p, q \leq s$ such that $X = X_p$ and $Y = X_q$. In $M(E_{n+1}, C^{n+1})$, we have

$$\mathbf{s}(X_p) = \sum_{x_p \in X_p} s(x_p) = \sum_{(x_1, \dots, x_s) \in \prod_{i=1}^s X_i} v(x_1, \dots, x_s) = \sum_{x_q \in X_q} s(x_q) = \mathbf{s}(X_q). \tag{4.2}$$

Hence $\mathbf{s}(X) = \mathbf{s}(Y)$ for all $X, Y \in C_v$, which shows that ψ_n is well-defined. Clearly φ_n and ψ_n are mutually inverse, so we get that φ_n is an isomorphism.

(b) By Lemma 3.8, the natural map $M(\iota_{V_n}) \colon M(E_n, C^n) \to M(E_{V_n}, C^{V_n})$ is a unitary embedding. Hence the map $\iota_n = \varphi_n^{-1} \circ M(\iota_{V_n}) \colon M(E_n, C^n) \to M(E_{n+1}, C^{n+1})$ is a unitary embedding.

(c) We use induction on n. For $n = 0$, we have $(E_0, C^0) = (F_0, D^0)$. Assume that, for some $n \geq 0$, the natural map $j_n \colon (E_n, C^n) \to (F_n, D^n)$ induces an isomorphism $M(j_n) \colon M(E_n, C^n) \to M(F_n, D^n)$. Since $(F_{n+1}, D^{n+1}) = ((F_n)_{V_n}, (D^n)^{V_n})$, we get an isomorphism $M(E_{V_n}, C^{V_n}) \to M(F_{n+1}, D^{n+1})$ extending $M(j_n)$. Now we have the following commutative diagram:

$$\begin{tikzcd}
M(E_n, C^n) \arrow[r, "M(\iota_{V_n})"] \arrow[d, "M(j_n)"'] & M(E_{V_n}, C^{V_n}) \arrow[d, "\cong"'] & M(E_{n+1}, C^{n+1}) \arrow[l, "\varphi_n"'] \arrow[d, "M(j_{n+1})"] \\
M(F_n, D^n) \arrow[r] & M(F_{n+1}, D^{n+1}) \arrow[r, "\mathrm{id}"] & M(F_{n+1}, D^{n+1})
\end{tikzcd} \tag{4.3}$$

Since φ_n is an isomorphism by (a), we get that $M(j_{n+1})$ is also an isomorphism.

(d) Write $\mathfrak{R} = \varinjlim (M(\iota_n), \iota_n)$. We have from (c) a commutative diagram

$$\begin{tikzcd}
M(E_0, C^0) \arrow[r, "\iota_0"] \arrow[d, "\mathrm{id}"'] & M(E_1, C^1) \arrow[r, "\iota_1"] \arrow[d, "M(j_1)"', "\cong"] & M(E_2, C^2) \arrow[r, dotted] \arrow[d, "M(j_2)"', "\cong"] & \mathfrak{R} \arrow[d] \\
M(F_0, D^0) \arrow[r] & M(F_1, D^1) \arrow[r] & M(F_2, D^2) \arrow[r, dotted] & M(F_\infty, D^\infty)
\end{tikzcd}$$

such that all the vertical maps $M(j_n)\colon M(E_n, C^n) \to M(F_n, D^n)$ are isomorphisms, so we get an isomorphism $\mathfrak{R} \to M(F_\infty, D^\infty)$. This shows the result. $\square$

Given a finite presentation $\langle \mathcal{X} \mid \mathcal{R} \rangle$ of a finitely presented abelian conical monoid M as in Definition 4.2, we can associated to it the refinement monoid

$$\mathfrak{R}(\mathcal{X} \mid \mathcal{R}) = M(F_\infty, D^\infty)\,,$$

where (F_∞, D^∞) is the complete multiresolution of the bipartite separated graph associated to the presentation $\langle \mathcal{X} \mid \mathcal{R} \rangle$. Observe that this construction depends strongly on the presentation, not just the monoid M. For instance, we may consider two different presentations $\langle a, b \mid a = a, b = b \rangle$ and $\langle a, b \mid a + b = a + b \rangle$ of the free abelian monoid on two generators F. In the first presentation the resulting $\mathfrak{R}$ is just F. For the second presentation, the corresponding refinement monoid $\mathfrak{R}$ is atomless, and is isomorphic to the monoid $\mathcal{V}(K[G])$, where G is the lamplighter group, see Example 9.7.

If we consider the standard presentation

$$M(E) = \langle a_v \mid a_v = \sum_{e \in r^{-1}(v)} a_{s(e)} \quad (v \in E^0) \rangle$$

of a graph monoid $M(E)$ associated to a non-separated finite graph E, then it can be shown that $\mathfrak{R} = M(E)$. Since $\mathfrak{R}$ is a refinement monoid, we obtain an alternative proof of [10, Proposition 4.4] for finite graphs.

5. The main construction

This section contains our main construction on *algebras*. Let (E, C) be a finite bipartite separated graph, with $r(E^1) = E^{0,0}$ and $s(E^1) = E^{0,1}$. Let $\{(E_n, C^n)\}_{n \geq 0}$ be the canonical sequence of bipartite separated graphs associated to it (see Definition 4.4(c)), and let B_n be the commutative, finite dimensional subalgebra of $L(E_n, C^n)$ generated by E_n^0. Here $L(E_n, C^n)$ stands for the Leavitt path *-algebra of the separated graph (E_n, C^n) over a fixed field with involution $(K, *)$.

Theorem 5.1. *With the above notation, for each $n \geq 0$, there exists a surjective $*$-algebra homomorphism*

$$\phi_n \colon L(E_n, C^n) \twoheadrightarrow L(E_{n+1}, C^{n+1}).$$

Moreover, the following properties hold:

(a) $\ker(\phi_n)$ *is the ideal I_n of $L(E_n, C^n)$ generated by all the commutators $[ee^*, ff^*]$, with $e, f \in E^1_n$, so that $L(E_{n+1}, C^{n+1}) \cong L(E_n, C^n))/I_n$.*

(b) *The restriction of ϕ_n to B_n defines an injective homomorphism from B_n into B_{n+1}.*

(c) *There is a commutative diagram*

$$\begin{tikzcd} M(E_n, C^n) \arrow[r, "\iota_n"] \arrow[d, "\cong"'] & M(E_{n+1}, C^{n+1}) \arrow[d, "\cong"] \\ \mathcal{V}(L(E_n, C^n)) \arrow[r, "\mathcal{V}(\phi_n)"] & \mathcal{V}(L(E_{n+1}, C^{n+1})) \end{tikzcd} \tag{5.1}$$

where the vertical maps are the canonical maps, which are isomorphisms by [8, Theorem 4.3].

Proof. Set $A_n = L(E_n, C^n)$. Define $\phi_n \colon A_n \to A_{n+1}$ on vertices $u \in E^{0,0}_n$ by the formula

$$\phi_n(u) = \sum_{(x_1,\dots,x_{k_u}) \in \prod_{i=1}^{k_u} X^u_i} v(x_1, \dots, x_{k_u}),$$

where $C_u = \{X_1, \dots, X_{k_u}\}$, and by $\phi_n(w) = w$ for all $w \in E^{0,1}_n$. For an arrow $x_i \in X^u_i$, define

$$\phi_n(x_i) = \sum_{x_j \in X^u_j, j \neq i} \left(\alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u})\right)^*,$$

where $\alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u}) = \alpha^{x_i}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_{k_u})$. We have to show that the defining relations of $L(E_n, C^n)$ are preserved by ϕ_n. It is easily checked that (V) and (E) are preserved by ϕ_n. To see that (SCK1) is preserved by ϕ_n, take $u \in E^{0,0}_n$ and $x_i, x'_i \in X^u_i$ for some i. Observe that

$$\phi_n(x_i)^* \phi_n(x'_i) = \sum_{x_j, y_j \in X^u_j, j \neq i} \alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u}) \alpha^{x'_i}(y_1, \dots, \widehat{x'_i}, \dots, y_{k_u})^*.$$

If $x_i \neq x'_i$, then $v(x_1, \dots, x_i, \dots, x_{k_u}) v(y_1, \dots, x'_i, \dots, y_{k_u}) = 0$ for all $x_j, y_j \in X^u_j$, $j \neq i$, and so $\phi_n(x_i)^* \phi_n(x'_i) = 0$. If $x_i = x'_i$ then $v(x_1, \dots, x_i, \dots, x_{k_u}) v(y_1, \dots, x_i, \dots, y_{k_u}) = 0$ except when $x_j = y_j$ for all $j \neq i$, and, hence recalling that $X(x_i) \in C^{n+1}_{s(x_i)}$, we get

$$\phi_n(x_i)^* \phi_n(x'_i) = \sum_{x_j \in X^u_j, j \neq i} \alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u}) \alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u})^* = \sum_{z \in X(x_i)} zz^* = s(x_i),$$

as desired. Finally we check that (SCK2) is also preserved by ϕ_n. Take $u \in E_n^{0,0}$ and $X_i^u \in C_u^n$. Then

$$\begin{aligned}\sum_{x_i \in X_i^u} \phi_n(x_i)\phi_n(x_i)^* &= \sum_{x_i \in X_i^u} \sum_{x_j \in X_j^u, j \neq i} \alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u})^* \alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u}) \\ &= \sum_{(x_1, \dots, x_{k_u}) \in \prod_{j=1}^{k_u} X_j^u} v(x_1, \dots, x_{k_u}) = \phi_n(u)\,,\end{aligned}$$

as wanted.

To check surjectivity, it is enough to check that all vertices and edges in E_{n+1} belong to the image of ϕ_n. Clearly, $w \in \phi_n(A_n)$ for all $w \in E_{n+1}^{0,0}$. Now take $x_i \in X_i^u$, for $i = 1, 2, \dots, k_u$. We have

$$\phi_n(x_i)\phi_n(x_i)^* = \sum_{y_j \in X_j^u, j \neq i} v(y_1, \dots, x_i, \dots, y_{k_u})\,. \tag{5.2}$$

Therefore, we get

$$v(x_1, x_2, \dots, x_{k_u}) = \phi_n((x_1x_1^*)(x_2x_2^*)\cdots(x_{k_u}x_{k_u}^*))\,, \tag{5.3}$$

showing that $v(x_1, x_2, \dots, x_n) \in \phi_n(A_n)$. Moreover, we get

$$\alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u})^* = v(x_1, \dots, x_{k_u})\phi_n(x_i)\,, \tag{5.4}$$

which, together with (5.3) gives that $\alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u}) \in \phi_n(A_n)$. This concludes the proof that ϕ_n is surjective.

We now proceed to show (a), (b), (c).

(a) It follows from (5.2) that $\{\phi_n(ee^*) \mid e \in E_n^1\}$ is a family of commuting projections in $L(E_{n+1}, C^{n+1})$. Therefore all commutators $[ee^*, ff^*]$ are contained in the kernel of ϕ_n, and we obtain an induced surjective map $\overline{\phi}_n \colon A_n/I_n \to A_{n+1}$. We define a map $\gamma_n \colon A_{n+1} \to A_n/I_n$ by $\gamma_n(w) = w$ for $w \in E_{n+1}^{0,0}$, and

$$\gamma_n(v(x_1, \dots, x_{k_u})) = (x_1x_1^*)(x_2x_2^*)\cdots(x_{k_u}x_{k_u}^*),$$

$$\gamma_n(\alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u})) = x_i^*(x_1x_1^*)(x_2x_2^*)\cdots(x_{k_u}x_{k_u}^*)$$

for $(x_j) \in \prod_{j=1}^{k_u} X_j$. Using the commutativity in A_n/I_n of the set of projections $\{ee^* \mid e \in E_n^1\}$, and the defining relations of $L(E_n, C^n)$, it is easy to check that the defining relations of $A_{n+1} = L(E_{n+1}, C^{n+1})$ are preserved by γ_n. Clearly, γ_n is the inverse of $\overline{\phi}_n$, and so we get that $\ker(\phi_n) = I_n$.

(b) This follows from the definition of ϕ_n.

(c) Denote by $\tau_n \colon M(E_n, C^n) \to \mathcal{V}(A_n)$ the natural map sending a_v to $[v]$, for $v \in E_n^0$. This map is an isomorphism by [8, Theorem 4.3]. Recall that $\iota_n = \varphi_n^{-1} \circ M(\iota_{V_n}) = \psi_n \circ M(\iota_{V_n})$ has been defined in the proof of Lemma 4.5(b).

If $w \in E_n^{0,1}$ then $(\tau_{n+1} \circ \iota_n)(a_w) = [w] = (\mathcal{V}(\phi_n) \circ \tau_n)(a_w)$. If $u \in E_n^{0,0}$, then it follows from the definition of ψ_n, formula (4.2) and the definition of ϕ_n that

$$(\tau_{n+1} \circ \iota_n)(a_u) = \sum_{(x_j) \in \prod_{j=1}^{k_u} X_j^u} [v(x_1, \dots, x_{k_u})] = (\mathcal{V}(\phi_n) \circ \tau_n)(a_u)\,,$$

so that we get the commutativity of the diagram (5.1). $\square$

We now study in more detail the relationship between the different layers. To simplify the notation, we will write $D_n = F^{0,n} = E_n^{0,0}$ for all $n \geq 0$.

Note that, for $n \geq 2$ we have a surjective map $r_n \colon D_n \to D_{n-2}$ given by $r_n(v(x_1, \dots x_{k_u})) = u$, where $u \in D_{n-2}$ and $x_i \in X_i^u$, and where, as usual, $C_u = \{X_1^u, \dots, X_{k_u}^u\}$. For $n = 2m$, we thus obtain a surjective map $\mathfrak{r}_{2m} = r_2 \circ r_4 \circ \cdots \circ r_{2m} \colon D_{2m} \to D_0$. Similarly, we have a map $\mathfrak{r}_{2m+1} = r_3 \circ r_5 \circ \cdots \circ r_{2m+1} \colon D_{2m+1} \to D_1$. We call $\mathfrak{r}(v)$ the *root* of v. Observe that we have

$$D_{2n} = \bigsqcup_{v \in D_0} \mathfrak{r}_{2n}^{-1}(v); \qquad D_{2n+1} = \bigsqcup_{v \in D_1} \mathfrak{r}_{2n+1}^{-1}(v) \tag{5.5}$$

Lemma 5.2. (1) *For all $n \geq 1$ and $v \in D_{2n}$ we have $|s^{-1}(v)| = |C_v| = |C_{\mathfrak{r}_{2n}(v)}|$.*

(2) *For all $n \geq 0$ and $w \in D_{2n+1}$ we have $|s^{-1}(w)| = |C_w| = |C_{\mathfrak{r}_{2n+1}(w)}|$.*

Proof. (1) By induction, it suffices to show that $|s^{-1}(v)| = |C_v| = |C_{r_{2n}(v)}|$ for every $v \in D_{2n}$, $n \geq 1$. Let $v \in D_{2n}$, $n \geq 1$, and write $v = v(x_1, \dots, x_{k_u})$, where $x_i \in X_i^u$ and $C_u = \{X_1^u, \dots, X_{k_u}^u\}$. Note that $s^{-1}(v)$ has exactly k_u elements, namely

$$\alpha^{x_1}(x_2, \dots, x_{k_u}), \dots, \alpha^{x_{k_u}}(x_1, \dots, x_{k_u - 1}).$$

Thus $|s^{-1}(v)| = k_u = |C_u| = |C_{r_{2n}(v)}|$. On the other hand, by definition $C_v = \{X(y) \mid y \in s_{E_{2n-1}}^{-1}(v)\}$, so that $|C_v| = |s^{-1}(v)| = |C_u|$, as desired.

(2) This is proved exactly as in (1). $\square$

Remark 5.3. Fix an enumeration $\{X_1^u, \dots, X_{k_u}^u\}$ for the elements of C_u, where $u \in D_0 = E^{0,0}$. This gives an enumerations of the elements in $s^{-1}(v)$ for $v \in r_2^{-1}(u)$. Indeed, for such an element v we have $v = v(x_1, \dots, x_{k_n})$, with $x_i \in X_i^u$, and we can set $z_i = \alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u})$, so that we obtain the enumeration $\{z_1, \dots, z_{k_u}\}$ of $s^{-1}(v)$. This in turn gives an enumeration of the elements of the set C_v, namely $C_v = \{X_1^v, \dots, X_{k_u}^v\}$, where $X_i^v = X(z_i)$ for all i. Obviously this translates to all the vertices $v \in D_{2n}$, so that we obtain canonical enumerations of the elements of $s^{-1}(v)$ and of the elements of C_v.

Define $\Phi_n = \phi_{n-1} \circ \cdots \circ \phi_0 \colon L(E, C) \to L(E_n, C^n)$.

Lemma 5.4. *Let $u \in D_0$ and set $C_u = \{X_1^u, \dots X_{k_u}^u\}$. For each $n \geq 1$, there exists a partition*

$$\mathfrak{r}_{2n}^{-1}(u) = \bigsqcup_{x_i \in X_i^u} Z_{2n}(x_i)\,,$$

such that, for each $x_i \in X_i^u$*, we have*

$$\Phi_{2n}(x_i) = \sum_{v\in Z_{2n}(x_i)} \sum_{x\in X_i^v} x\,, \tag{5.6}$$

where the enumeration of the elements in C_v*,* $v \in D_{2n}$*, is the canonical one described in Remark 5.3.*

Proof. We first prove the case $n = 1$. Observe that $\phi_0(x_i) = \sum_{y\in X(x_i)} y^*$. The elements $y \in X(x_i)$ are of the form $\alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u})$, and we set

$$Z_2(x_i) = \{s(y) \mid y \in X(x_i)\} = \{v(x_1, \dots, x_i, \dots, x_{k_u}) \mid x_j \in X_j^u, j \neq i\}.$$

Note that $s(y) \neq s(y')$ for $y, y' \in X(x_i)$ with $y \neq y'$. Observe that we get a partition

$$\mathfrak{r}_2^{-1}(u) = \bigsqcup_{x_i\in X_i^u} Z_2(x_i).$$

Let $v = v(x_1, \dots, x_i, \dots, x_{k_u}) \in Z_2(x_i)$. Then in the canonical enumeration of the elements of $s^{-1}(v)$, say $\{z_1, \dots, z_{k_u}\}$, the element $\alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u})$ corresponds to z_i, and there is no other element of $X(x_i)$ having source equal to v. The corresponding element $X(y) = X(z_i)$ of C_v is the element labelled as X_i^v (see Remark 5.3). Therefore we get

$$\Phi_2(x_i) = \phi_1\Big(\sum_{y\in X(x_i)} y^*\Big) = \sum_{y\in X(x_i)} \sum_{x\in X(y)} x = \sum_{v\in Z_2(x_i)} \sum_{x\in X_i^v} x\,.$$

Assume now that $n \geq 1$ and that (5.6) holds. Note that

$$\phi_{2n}(\Phi_{2n}(x_i)) = \sum_{v\in Z_{2n}(x_i)} \sum_{x_i'\in X_i^v} \sum_{y\in X(x_i')} y^*. \tag{5.7}$$

Observe that $\bigsqcup_{x_i'\in X_i^v}\{s(y) \mid y \in X(x_i')\} = r_{2n+2}^{-1}(v)$, and that $s(y) \neq s(y')$ for $y \neq y'$ and $y, y' \in \bigcup_{v\in Z_{2n}(x_i)} \bigcup_{x_i'\in X_i^v} X(x_i')$. Set

$$Z_{2n+2}(x_i) = \bigsqcup_{v\in Z_{2n}(x_i)} r_{2n+2}^{-1}(v) = r_{2n+2}^{-1}(Z_{2n}(x_i)).$$

Clearly we get a partition $\mathfrak{r}_{2n+2}^{-1}(u) = \bigsqcup_{x_i\in X_i^u} Z_{2n+2}(x_i)$. Moreover, arguing as in the case $n = 1$ we get

$$\begin{aligned}\Phi_{2n+2}(x_i) &= \phi_{2n+1}\Big(\sum_{v\in Z_{2n}(x_i)} \sum_{x_i'\in X_i^v} \sum_{y\in X(x_i')} y^*\Big)\\ &= \sum_{v\in Z_{2n}(x_i)} \sum_{x_i'\in X_i^v} \sum_{y\in X(x_i')} \sum_{x\in X(y)} x = \sum_{v'\in Z_{2n+2}(x_i)} \sum_{x\in X_i^{v'}} x\,,\end{aligned}$$

completing the induction step. $\square$

Lemma 5.5. *Let $u \in D_0$ and set $C_u = \{X^u_1, \dots X^u_{k_u}\}$. For $x_i \in X^u_i$, $n \geq 0$ and $b \in B_n$ we have*

$$\Phi_{n+1}(x_i)\phi_n(b)\Phi_{n+1}(x_i)^* \in B_{n+1}, \qquad \Phi_{n+1}(x_i)^*\phi_n(b)\Phi_{n+1}(x_i) \in B_{n+1}.$$

Moreover, if b is a projection in B_n, then both $\Phi_{n+1}(x_i)\phi_n(b)\Phi_{n+1}(x_i)^*$ and $\Phi_{n+1}(x_i)^*\phi_n(b)\Phi_{n+1}(x_i)$ are projections in B_{n+1}.

Proof. Note that it is enough to check that $\phi_n(\Phi_n(x_i)w\Phi_n(x_i)^*) \in B_{n+1}$ and $\phi_n(\Phi_n(x_i)^*w\Phi_n(x_i)) \in B_{n+1}$ for all vertices $w \in E^0_n$.

We start with the cases $n = 0$ and $n = 1$. For $n = 0$ the result is clear by (SCK1) and (5.2). For $n = 1$, observe that $\phi_0(x_i)w\phi_0(x_i)^*$ is 0 except when $w = s(x_i)$ and in this case $\phi_0(x_i)w\phi(x_i)^* = \phi_0(x_ix_i^*) \in B_1$ by (5.2). On the other hand, $\phi_0(x_i)^*w\phi_0(x_i)$ is 0 except for $v = v(x_1, \dots, x_i, \dots x_{k_u})$ for some $x_j \in X^u_j$, $j \neq i$. In this case we have

$$\phi_0(x_i)^*v(x_1, \dots, x_{k_u})\phi_0(x_i) = \alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u})\alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u})^*\,,$$

so that $\phi_1(\phi_0(x_i)^*v(x_1, \dots, x_{k_u})\phi_0(x_i)) \in B_2$ again by (5.2).

Now we will show the result for an even index $2n$, with $n \geq 1$. By (5.6), we have

$$\Phi_{2n}(x_i)w\Phi_{2n}(x_i)^* = (\sum_{v\in Z_{2n}(x_i)}\sum_{x\in X^v_i} x)w(\sum_{v\in Z_{2n}(x_i)}\sum_{x\in X^v_i} x)^*.$$

Since $n \geq 1$, the projections $\{s(x) \mid x \in \bigcup_{v\in Z_{2n}(x_i)} X^v_i\}$ are pairwise orthogonal, and we conclude that the above term is 0 except when $w = s(x)$ for a, necessarily unique, $x \in \bigcup_{v\in Z_{2n}(x_i)} X^v_i$. In this case ,we get

$$\Phi_{2n}(x_i)w\Phi_{2n}(x_i)^* = xx^*,$$

and so $\phi_{2n}(\Phi_{2n}(x_i)w\Phi_{2n}(x_i)^*) = \phi_{2n}(xx^*) \in B_{2n+1}$ by (5.2).

On the other hand, note that $\Phi_{2n}(x_i)^*v\Phi_{2n}(x_i) = 0$ except for $v \in Z_{2n}(x_i)$. If $v \in Z_{2n}(x_i)$, the we get

$$\Phi_{2n}(x_i)^*v\Phi_{2n}(x_i) = (\sum_{x\in X^v_i} x^*)(\sum_{y\in X^v_i} y) = \sum_{x\in X^v_i} x^*x = \sum_{x\in X^v_i} s(x) \in B_{2n}\,.$$

We conclude that $\Phi_{2n}(x_i)$ is a partial isometry in A_{2n} with initial projection $\Phi_{2n}(x_i)^*\Phi_{2n}(x_i) = \sum_{v\in Z_{2n}(x_i)}\sum_{x\in X^v_i} s(x)$ and final projection $\Phi_{2n}(x_i)\Phi_{2n}(x_i)^* = \sum_{v\in Z_{2n}(x_i)} v$. This shows the result for the index $2n$.

For an odd integer $2n+1$, with $n \geq 1$, use (5.7) and an argument similar to the above to conclude the result. In this case $\Phi_{2n+1}(x_i)$ is a partial isometry in A_{2n+1} with initial projection $\Phi_{2n+1}(x_i)^*\Phi_{2n+1}(x_i) = \sum_{v\in Z_{2n}(x_i)}\sum_{x\in X^v_i} s(x)$ and final projection $\Phi_{2n+1}(x_i)\Phi_{2n+1}(x_i)^* = \sum_{v\in Z_{2n+2}(x_i)} v$. □

Theorem 5.6. *Let U be the subsemigroup of the multiplicative semigroup of $A_0 = L(E, C)$ generated by $\{x, x^* : x \in E^1\}$. For $w \in U$ set $e(w) = ww^*$. For $n \geq 1$, let J_n be the ideal of A_0 generated by all the commutators $[e(w), e(w')]$, with w, w' elements of U which are products of $\leq n$ generators $\{x, x^* : x \in E^1\}$. Then* $\ker(\Phi_n) = J_n$ *so that $A_n \cong A_0/J_n$.*

Proof. Observe that, for $n = 1$ this follows from Theorem 5.1(a).

Write U_n for the set of elements in U which can be written as a product of $\leq n$ generators. Using Lemma 5.5 it is very simple to show inductively that $\Phi_n(e(w)) = \Phi_n(w)\Phi_n(w)^*$ is a projection in B_n for any $w \in U_n$. Since B_n is a commutative algebra, we deduce that $[e(w), e(w')] \in \ker(\Phi_n)$ if $w, w' \in U$ are products of $\leq n$ generators. This shows that $J_n \subseteq \ker(\Phi_n)$ for all $n \geq 1$.

Before we prove the reverse inclusion, we need to establish a claim.

Claim: (1) If $z \in E^1_{2n}$, then there exists $x \in E^1$ and $w_1, \dots w_t \in U_{2n}$ such that $\Phi_{2n}(e(w_j)) \leq \Phi_{2n}(x^*x)$ and

$$z = \Phi_{2n}(xe(w_1)e(w_2)\cdots e(w_t)). \tag{5.8}$$

(2) If $z \in E^1_{2n+1}$, then there exists $x \in E^1$ and $w_1, \dots w_t \in U_{2n+1}$ such that $\Phi_{2n+1}(e(w_j)) \leq \Phi_{2n}(xx^*)$ and

$$z = \Phi_{2n+1}(x^*e(w_1)e(w_2)\cdots e(w_t)). \tag{5.9}$$

Proof of Claim. The claim is proved by induction. It trivially holds for $z \in E^1_0$, and it holds for $z \in E^1_1$ by (5.4) and (5.3). Assume that (1) holds for all $z \in E^1_{2n}$, for some $n \geq 0$: Then we will show that (2) holds for all $z \in E^1_{2n+1}$. So, let $z \in E^1_{2n+1}$. Then there exists $x_j \in X^u_j$, $j = 1, \dots, k_u$, for some $u \in D_{2n} = E^{0,0}_{2n}$, such that $z = \alpha^{x_i}(x_1, \dots, \widehat{x_i}, \dots, x_{k_u})$, and it follows from (5.4) that

$$z = \phi_{2n}(x_i^*)v(x_1, \dots, x_{k_u}).$$

By induction hypothesis, we have that there is $x \in E^1$ and $w_1, \dots, w_t \in U_{2n}$ such that $\Phi_{2n}(e(w_j)) \leq \Phi_{2n}(x^*x)$ and $x_i = \Phi_{2n}(xe(w_1)\dots e(w_t))$. It follows that

$$z = \Phi_{2n+1}(xe(w_1)\cdots e(w_t))^*v(x_1, \dots, x_{k_u}). \tag{5.10}$$

But now it follows from Lemma 5.5 that

$$\Phi_{2n+1}(xe(w_1)\cdots e(w_t)) = \Phi_{2n+1}(e(xw_1)\cdots e(xw_t)x). \tag{5.11}$$

On the other hand, using again Lemma 5.5, the induction hypothesis and (5.3), we see that there are $w'_1, \dots w'_s$ in U_{2n+1} such that

$$v(x_1, \dots, x_{k_u}) = \Phi_{2n+1}(e(w'_1)\cdots e(w'_s)). \tag{5.12}$$

Substituting (5.11) and (5.12) in (5.10) we get

$$z = \Phi_{2n+1}(x^*e(xw_t)\cdots e(xw_1)e(w'_1)\cdots e(w'_s))\,,$$

showing (5.9). A similar argument shows that (2) implies (1) for the index $2n + 2$. This concludes the proof of the Claim.

Now we follow with the proof of the theorem. We show the reverse inclusion $\ker(\Phi_m) \subseteq J_m$ by induction on m. So, assume that the result holds for some $m \geq 1$ and let us show that it also holds for $m + 1$. Observe that, by Theorem 5.1(a), $\ker\Phi_{m+1}/\ker\Phi_m$ is the ideal of A_m generated by $[zz^*, (z')(z')^*]$, where $z, z' \in E^1_m$. Assume first that $m = 2n$ for some $n \geq 0$ and that $z, z' \in E^1_{2n}$. By the first part of the Claim, there exist then $x, x' \in E^1$ and

$w_1, \dots w_t, w'_1, \dots, w'_{t'} \in U_{2n}$ such that $\Phi_{2n}(e(w_j)) \leq \Phi_{2n}(x^*x)$, $\Phi_{2n}(e(w'_j)) \leq \Phi_{2n}((x')^*(x'))$, and

$$z = \Phi_{2n}(xe(w_1)e(w_2)\cdots e(w_t)), \qquad z' = \Phi_{2n}(x'e(w'_1)e(w'_2)\cdots e(w'_{t'})).$$

Since $\ker \Phi_{2n} = J_{2n}$ by induction hypothesis, and $J_{2n} \subseteq J_{2n+1}$, it is enough to check that

$$\Big[xe(w_1)\cdots e(w_t)e(w_t)^* \cdots e(w_1)^*x^* \, , \, x'e(w'_1)\cdots e(w'_{t'})e(w'_{t'})^* \cdots e(w'_1)^*(x')^*\Big] \in J_{2n+1}.$$

We have

$$\begin{aligned}
&\Big[xe(w_1)\cdots e(w_t)e(w_t)^* \cdots e(w_1)^*x^* + J_{2n} \, , \, x'e(w'_1)\cdots e(w'_{t'})e(w'_{t'})^* \cdots e(w'_1)^*(x')^* + J_{2n}\Big] \\
&= \Big[xe(w_1)\cdots e(w_t)x^* + J_{2n} \, , \, x'e(w'_1)\cdots e(w'_{t'})(x')^* + J_{2n}\Big] \\
&= \Big[xe(w_1)(x^*x)e(w_2)\cdots (x^*x)e(w_t)x^* + J_{2n} \, , \, x'e(w'_1)(x')^*x'e(w'_2)\cdots (x')^*x'e(w'_{t'})(x')^* + J_{2n}\Big] \\
&= \Big[e(xw_1)e(xw_2)\cdots e(xw_t) + J_{2n} \, , \, e(x'w'_1)e(x'w'_2)\cdots e(x'w'_{t'}) + J_{2n}\Big] \\
&\subseteq \sum_{i,j} A_0[e(xw_i), e(x'w'_j)]A_0 + J_{2n} \subseteq J_{2n+1} \, ,
\end{aligned}$$

where we have used Leibnitz rule for the first containment. A similar argument, using part (2) of the Claim instead of part (1), gives the proof of the inductive step for $m = 2n+1$. This completes the proof of the theorem. □

Write $A_\infty = \varinjlim A_n$,and $B_\infty = \varinjlim B_n$. We have a commutative diagram as follows:

$$\begin{tikzcd}
B_0 \arrow[r] \arrow[d] & B_1 \arrow[r] \arrow[d] & B_2 \arrow[r, dotted] \arrow[d] & B_\infty \arrow[d] \\
A_0 \arrow[r, "\phi_0"] & A_1 \arrow[r, "\phi_1"] & A_2 \arrow[r, dotted] & A_\infty
\end{tikzcd}$$

All the maps $B_n \to B_{n+1}$ are injective and all the maps $A_n \to A_{n+1}$ are surjective.

Notation 5.7. We write $L^{\mathrm{ab}}(E,C) = A_\infty$. Observe that, by Theorem 5.6, $L^{\mathrm{ab}}(E,C) \cong L(E,C)/J$, where $J = \bigcup_{n=1}^\infty J_n$ is the ideal generated by all the commutators $[e(w), e(w')]$, with $w, w' \in U$. If we use in the above construction the full graph C*-algebras $C^*(E_n, C^n)$ instead of the Leavitt path algebras $L(E_n, C^n)$, we obtain in the limit a certain C*-algebra, that we denote by $\mathcal{O}(E,C)$, in analogy with the notation introduced in [5].

Corollary 5.8. *We have* $\mathcal{V}(L^{\mathrm{ab}}(E,C)) \cong \varinjlim(M(E_n, C^n), \iota_n) \cong M(F_\infty, D^\infty)$. *Consequently the natural map* $M(E,C) \to \mathcal{V}(L^{\mathrm{ab}}(E,C))$ *is a refinement of* $M(E,C)$.

Proof. This follows immediately from the continuity of the functor $\mathcal{V}$, Theorem 5.1(c) and Lemma 4.5(d). $\square$

Remark 5.9. Assume that $K = \mathbb{C}$, the complex field. Then the commutative $*$-algebra B_∞ is an algebraic direct limit of finite-dimensional commutative C*-algebras B_n, with injective maps $\phi_n|_{B_n} \colon B_n \to B_{n+1}$. Since each B_n sits as a C^*-subalgebra of $C^*(E_n, C^n)$ (cf. [7, Theorem 3.8(1)]), it follows that the C^*-completion $\mathcal{B}_0$ of B_∞ sits as a C*-subalgebra of $\mathcal{O}(E, C)$.

Recall the following definition from [23]. (The definition has been conveniently modified to work in the purely algebraic situation as well as in the C*-context.)

Definition 5.10. A set S of partial isometries in a $*$-algebra $\mathcal{A}$ is said to be *tame* if every element of $U = \langle S \cup S^* \rangle$, the multiplicative semigroup generated by $S \cup S^*$, is a partial isometry, and the set $\{e(u) \mid u \in U\}$ of final projections of the elements of U is a commuting set of projections of $\mathcal{A}$.

We are now ready to show that $L^{\mathrm{ab}}(E, C)$ is the universal algebra generated by a *tame* set of partial isometries $\underline{x}$, for $x \in E^1$ satisfying the defining relations of $L(E, C)$, which can be re-stated completely in terms of the elements in $E^1 \cup (E^1)^*$, indeed in terms of the final projections $e(u)$ of these elements, as follows:

(PI1) $e(x)e(y) = \delta_{x,y}e(x)$, for $x, y \in X$, $X \in C$.

(PI2) $e(x^*) = e(y^*)$ if $x, y \in E^1$ and $s(x) = s(y)$.

(PI3) $\sum_{x \in X} e(x) = \sum_{y \in Y} e(y)$ for all $X, Y \in C_v$, $v \in E^{0,0}$.

For $w \in E^{0,1}$ and $x \in E^1$ with $s(x) = w$, denote the projection $e(x^*)$ by p_w. Similarly, for $v \in E^{0,0}$, denote by p_v the projection corresponding to $\sum_{x \in X} e(x)$, for $X \in C_v$. (Note that these projections are well-defined by (PI1)–(PI3).)

(PI4) $p_v p_{v'} = 0$ for every pair v, v' of different vertices in E^0, and $\sum_{v \in E^0} p_v = 1$.

Note that the condition $\sum_{v \in E^0} p_v = 1$ can be deduced from the other conditions, because E is a finite graph. It is easily seen that for a finite bipartite separated graph satisfying our standing assumptions that $s(E^1) = E^{0,1}$ and $r(E^1) = E^{0,0}$, the algebra $L(E, C)$ is just the universal algebra generated by a set of partial isometries $x \in E^1$ subject to the above relations (PI1)–(PI4). A similar statement holds for the full graph C*-algebra $C^*(E, C)$.

Theorem 5.11. *The algebra $L^{\mathrm{ab}}(E, C)$ (respectively, the C*-algebra $\mathcal{O}(E, C)$) is the universal $*$-algebra (respectively, universal C*-algebra) generated by a tame set of partial isometries $\{\underline{x} \mid x \in E^1\}$ satisfying the relations (PI1)–(PI4).*

Proof. Let U be the multiplicative semigroup of $L(E, C)$ generated by $E^1 \cup (E^1)^*$, and let U_n be the set of those elements of U which can be written as a product of $\leq n$ generators. Then $L^{\mathrm{ab}}(E, C) = L(E, C)/J$, where $J = \bigcup_{n=1}^\infty J_n$ and J_n is the ideal generated by the commutators $[e(w), e(w')]$, with $w, w' \in U_n$ (Theorem 5.6). If $w \in U_n$ then it was observed in the proof of Theorem 5.6 that $e(w)$ is a projection in $A_n = L(E, C)/J_n$. It is rather easy to show, by induction on n, that this implies that every element of U_n is a partial isometry in $L(E, C)/J_n$.

Denoting by $\underline{u}$ the image of $u \in U$ through the canonical projection $L(E,C) \to L^{\mathrm{ab}}(E,C)$, we thus have that all elements $\underline{u}$ are partial isometries and the final projections of these elements commute with each other. It follows that $\{\underline{x} \mid x \in E^1\}$ is a tame set of partial isometries in $L^{\mathrm{ab}}(E,C)$. Since J is the ideal generated by all the commutators $[e(w), e(w')]$, with $w, w' \in U$, it is clear that $L^{\mathrm{ab}}(E,C)$ is the universal algebra generated by a tame set of partial isometries satisfying conditions (PI1)–(PI4), as claimed.

The same proof works for $\mathcal{O}(E,C)$. □

By Stone duality, there is a zero-dimensional metrizable compact space $\Omega(E,C)$ such that the Boolean algebra of projections in B_∞ is isomorphic to the Boolean algebra of clopen subsets of $\Omega(E,C)$. A convenient way of describing this space is as follows. Consider the commutative AF-algebra $\mathcal{B}_0$, which is the C^*-completion of the locally matricial algebra B_∞ (taking $K = \mathbb{C}$). Then there is a unique zero-dimensional metrizable compact topological space $\Omega(E,C)$ such that $\mathcal{B}_0 \cong C(\Omega(E,C))$. Observe that we have a basis of clopen sets for the topology of $\Omega(E,C)$ which is in bijective correspondence with the vertices of the graph F_∞. Note that for any field K, the algebra B_∞ is the algebra $C_K(\Omega(E,C))$ of continuous functions $f \colon \Omega(E,C) \to K$, where K is endowed with the discrete topology.

We record for later use a fundamental property of the algebra B_∞.

Proposition 5.12. *The algebra B_∞ is generated by all the projections $e(w)$, for $w \in U$.*

Proof. The algebra B_∞ is generated by the images under the limit maps of the projections $v \in D_n$, for $n \geq 0$. Now if $v \in D_0$ then it follows from (SCK2) that v is a sum of projections $e(x)$, for $x \in X$, $X \in C_v$. Similarly, (SCK1) gives that the projections in D_1 are of the form $e(x^*)$. Suppose now that $v \in D_{n+1}$ with $n \geq 1$. Then $v = v(x_1, \dots, x_{k_u})$, where $x_i \in X_i^u$, $C_u = \{X_1^u, \dots, X_{k_u}^u\}$ and $u \in D_{n-1}$. Then it follows from either (5.8) or (5.9), and equation (5.3) that $v(x_1, \dots, x_{k_u})$ is a product of projections of the form $\Phi_n(e(w))$, where $w \in U_n$. □

6. Dynamical systems

Definition 6.1. Let G be a group with identity element 1. A *partial representation* of G on a unital K-algebra $\mathcal{A}$ is a map $\pi \colon G \to \mathcal{A}$ such that $\pi(1) = 1_{\mathcal{A}}$ and

$$\pi(g)\pi(h)\pi(h^{-1}) = \pi(gh)\pi(h^{-1}), \quad \pi(g^{-1})\pi(g)\pi(h) = \pi(g^{-1})\pi(gh)$$

for all $g, h \in G$. If G is a free group on a set X and $|g|$ denotes the length of $g \in G$ with respect to X, we say that a partial representation π is *semi-saturated* in case $\pi(ts) = \pi(t)\pi(s)$ for all $t, s \in G$ with $|ts| = |t| + |s|$.

Definition 6.2. Let G be a group and let $\mathcal{B}$ be an associative K-algebra (possibly without unit). A *partial action* of G on $\mathcal{B}$ consists of a family of two-sided ideals $\mathcal{D}_g$ of $\mathcal{B}$ ($g \in G$) and algebra isomorphisms $\alpha_g \colon \mathcal{D}_{g^{-1}} \to \mathcal{D}_g$ such that

(i) $\mathcal{D}_1 = \mathcal{B}$ and α_1 is the identity map on $\mathcal{B}$;

(ii) $\alpha_h(\mathcal{D}_{h^{-1}} \cap \mathcal{D}_g) = \mathcal{D}_h \cap \mathcal{D}_{hg}$;

(iii) $\alpha_g \circ \alpha_h(x) = \alpha_{gh}(x)$ for each $x \in \alpha_h^{-1}(\mathcal{D}_h \cap \mathcal{D}_{g^{-1}})$.

Let (E,C) be a finite bipartite separated graph, with $r(E^1)=E^{0,0}$ and $s(E^1)=E^{0,1}$, as in Definition 4.1. Throughout this section, we will denote by $\mathbb{F}$ the free group on the set E^1 of edges of E.

Proposition 6.3. *The algebra $L^{\mathrm{ab}}(E,C)$ is the universal $*$-algebra for semi-saturated partial representations of $\mathbb{F}$ satisfying the relations (PI1)-(PI4). That is, there is a semi-saturated partial representation $\pi\colon \mathbb{F}\to L^{\mathrm{ab}}(E,C)$ sending each $x\in E^1$ to the element $\underline{x}$ of $L^{\mathrm{ab}}(E,C)$, and, moreover given any semi-saturated partial representation ρ of $\mathbb{F}$ on a unital $*$-algebra B such that the partial isometries $\rho(x)$, $x\in E^1\sqcup (E^1)^*$, satisfy relations (PI1)–(PI4), there is a unique $*$-algebra homomorphism $\varphi\colon L^{\mathrm{ab}}(E,C)\to B$ such that $\rho=\varphi\circ\pi$.*

Proof. Since $\{\underline{x}\mid x\in E^1\}$ is a tame set of partial isometries in $L^{\mathrm{ab}}(E,C)$, it follows from [23, Proposition 5.4] that there exists a (unique) semi-saturated partial representation $\pi\colon\mathbb{F}\to L^{\mathrm{ab}}(E,C)$ such that $\pi(x)=\underline{x}$ for all $x\in E^1$.

The universality follows from Theorem 5.11, just as in [5, 2.4]. □

Let $\pi\colon G\to\mathcal{A}$ be a partial representation of a group G on a unital K-algebra $\mathcal{A}$. Then the elements $\varepsilon_g=\pi(g)\pi(g^{-1})$ $(g\in G)$ are commuting idempotents such that

$$\pi(g)\varepsilon_h=\varepsilon_{gh}\pi(g),\qquad \varepsilon_h\pi(g)=\pi(g)\varepsilon_{g^{-1}h}, \tag{6.1}$$

see [20, page 1944]. Let $\mathcal{B}$ be the subalgebra of $\mathcal{A}$ generated by all the ε_g $(g\in G)$, and for $g\in G$ set $\mathcal{D}_g=\varepsilon_g\mathcal{B}$. By [20, Lemma 6.5], the maps $\alpha^\pi_g\colon\mathcal{D}_{g^{-1}}\to\mathcal{D}_g$ $(g\in G)$ given by

$$\alpha^\pi_g(b)=\pi(g)b\pi(g)^*$$

are isomorphisms of K-algebras which determine a partial action α^π of G on $\mathcal{B}$.

In the case where $\mathcal{A}=L^{\mathrm{ab}}(E,C)$ and $\pi\colon\mathbb{F}\to L^{\mathrm{ab}}(E,C)$ is the partial representation described in Proposition 6.3, we set $\varepsilon_w=e(w)\in B_\infty$, and we observe that, by Proposition 5.12, the algebra $\mathcal{B}$ generated by all the projections $e(w)$ $(w\in\mathbb{F})$ is precisely the commutative algebra B_∞. It follows that $\mathcal{D}_w=B_\infty e(w)$, the ideal of B_∞ generated by $e(w)$.

As noted above, it follows that the maps

$$\phi_w\colon B_\infty e(w^*)\longrightarrow B_\infty e(w)$$

defined by $\phi_w(b)=\pi(w)b\pi(w)^*$, give a partial action of $\mathbb{F}$ on B_∞, namely $\phi=\alpha^\pi$.

We now recall the notion of a crossed product by a partial action, see e.g. [20, Definition 1.2].

Definition 6.4. Given a partial action α of a group G on a K-algebra $\mathcal{B}$, the *crossed product* $\mathcal{B}\rtimes_\alpha G$ is the set of all finite formal sums $\{\sum_{g\in G}a_g\delta_g\mid a_g\in\mathcal{D}_g\}$, where δ_g are symbols. Addition is defined in the obvious way, and multiplication is determined by $(a_g\delta_g)\cdot(b_h\delta_h)=\alpha_g(\alpha_{g^{-1}}(a_g)b_h)\delta_{gh}$.

Note that, given a partial action of G on a unital K-algebra $\mathcal{B}$, we have a partial representation $G\to\mathcal{B}\rtimes_\alpha G$, defined by $g\mapsto 1_g\delta_g$ (see [20, Lemma 6.2].

We need a further definition.

Definition 6.5. A *partial* (E,C)*-action* on a unital commutative $*$-algebra $\mathcal{B}$ consists of a collection of projections $\{e(u) \mid u \in E^1 \cup (E^1)^*\}$ satisfying the conditions (PI1)-(PI4), together with a collection of $*$-algebra isomorphisms $\alpha_x\colon e(x^*)\mathcal{B} \to e(x)\mathcal{B}$ for every $x \in E^1$.

If $\mathcal{B}_1$ and $\mathcal{B}_2$ are two unital commutative $*$-algebras with partial (E,C) actions α and β respectively, then an *equivariant homomorphism* from $\mathcal{B}_1$ to $\mathcal{B}_2$ is a $*$-algebra homomorphism $\psi\colon \mathcal{B}_1 \to \mathcal{B}_2$ such that $\psi(e_\alpha(u)) = e_\beta(u)$ for all $u \in (E^1) \cup (E^1)^*$ and such that the following diagram

$$\begin{tikzcd} e_\alpha(x^*)\mathcal{B}_1 \arrow[r, "\psi|"] \arrow[d, "\alpha_x"'] & e_\beta(x^*)\mathcal{B}_2 \arrow[d, "\beta_x"'] \\ e_\alpha(x)\mathcal{B}_1 \arrow[r, "\psi|"] & e_\beta(x)\mathcal{B}_2 \end{tikzcd} \tag{6.2}$$

commutes for all $x \in E^1$.

A commutative $*$-algebra $\mathcal{B}$ with a partial (E,C)-action α is *universal for partial* (E,C)*-actions* if given any commutative $*$-algebra $\mathcal{C}$ with a partial (E,C)-action β, there is a unique equivariant homomorphism $\psi\colon \mathcal{B} \to \mathcal{C}$

Lemma 6.6. *Any partial* (E,C)*-action on a unital commutative* $*$*-algebra* $\mathcal{B}$ *can be extended to a partial action* α *of* $\mathbb{F}$ *on* $\mathcal{B}$ *such that* $\mathcal{D}_g = e(g)\mathcal{B}$*, with* $e(g)$ *a projection in* $\mathcal{B}$ *for all* $g \in \mathbb{F}$*. Moreover, the corresponding partial representation* $\pi\colon \mathbb{F} \to \mathcal{B} \rtimes_\alpha \mathbb{F}$*, defined by* $\pi(g) = e(g)\delta_g$ *is semi-saturated.*

Proof. Let α be a partial (E,C)-action on a unital commutative $*$-algebra $\mathcal{B}$. Let $\mathbf{B}$ be the Boolean algebra of projections of $\mathcal{B}$, and let $\mathcal{I}(\mathbf{B})$ be the inverse semigroup of partially defined $*$-algebra isomorphisms with domain and range of the form $\mathbf{b}\mathcal{B}$, with $\mathbf{b} \in \mathbf{B}$. Then α can be extended to a map (also denoted by α) from $\mathbb{F}$ to $\mathcal{I}(\mathbf{B})$ by the rule

$$\alpha_{g_1g_2\cdots g_k} = \alpha_{g_1} \cdot \alpha_{g_2} \cdot \cdots \cdot \alpha_{g_k}\,,$$

where $g_1g_2\cdots g_k$ is the reduced form of an element of $\mathbb{F}$ and $\alpha_{x^{-1}} = \alpha_x^{-1}$ for all $x \in E^1$. It is easily checked that this extension provides a partial action on $\mathcal{B}$. See also [43, Example 2.4].

Finally we show that the representation $\pi\colon \mathbb{F} \to \mathcal{B} \rtimes_\alpha \mathbb{F}$ is semi-saturated. Let $g,h \in \mathbb{F}$ be such that $|gh| = |g| + |h|$. The identity $e(gh)\delta_{gh} = (e(g)\delta_g)(e(h)\delta_h)$ is equivalent to $e(gh) \le e(g)$. Since $|gh| = |g||h|$, it is evident from the definition that $e(gh) \le e(g)$, because $\alpha_{gh} = \alpha_g \cdot \alpha_h$ in $\mathcal{I}(\mathbf{B})$. □

Now we can obtain the basic relation between universality for partial representation obeying (PI1)-(PI4) and universality for partial (E,C) actions.

Theorem 6.7. *A partial representation* π *of* $\mathbb{F}$ *on* $\mathcal{A}$ *is universal for semi-saturated partial representations satisfying (PI1)-(PI4) if and only if* $\mathcal{A} \cong \mathcal{B} \rtimes_\alpha \mathbb{F}$*, where* $(\mathcal{B},\alpha)$ *is a universal partial* (E,C)*-action on the commutative* $*$*-algebra* $\mathcal{B}$*.*

Proof. Let π be a semi-saturated partial representation of $\mathbb{F}$ on the unital $*$-algebra $\mathcal{A}$ satisfying (PI1)-(PI4), and assume that $(\mathcal{A},\pi)$ is universal for semi-saturated partial representations of $\mathbb{F}$ satisfying (PI1)-(PI4). Let $\mathcal{B}$ be the commutative subalgebra of $\mathcal{A}$ generated by the projections $e(g) = \pi(g)\pi(g)^*$. Then $(\mathcal{B}, \alpha^\pi)$ is a partial (E, C)-action on $\mathcal{B}$, and $\mathcal{A} \cong \mathcal{B} \rtimes_{\alpha^\pi} \mathbb{F}$ by [20, Proposition 6.8] and the universal property of π. Assume that $(\mathcal{C}, \beta)$ is a partial (E, C)-action on a unital commutative $*$-algebra $\mathcal{C}$. By Lemma 6.6, β can be extended to a partial action, also denoted by β, of $\mathbb{F}$ on $\mathcal{C}$ such that $\mathcal{D}_g = e_\beta(g)\mathcal{C}$ for some projection $e_\beta(g)$, for all $g \in \mathbb{F}$, and such that the induced partial representation $\rho\colon \mathbb{F} \to \mathcal{C} \rtimes_\beta \mathbb{F}$ is semi-saturated. Since β is a partial (E, C)-action, the partial isometries $\rho(x)$, $x \in E^1$, satisfy the relations (PI1)-(PI4), so that there is a unique $*$-homomorphism $\varphi\colon \mathcal{B} \rtimes_\alpha \mathbb{F} \to \mathcal{C} \rtimes_\beta \mathbb{F}$ such that $\varphi(\pi(g)) = e_\beta(g)\delta_g$ for all $g \in \mathbb{F}$. In particular, we see that

$$\varphi(e(g)) = \varphi(\pi(g)\pi(g)^*) = (e_\beta(g)\delta_g)(e_\beta(g)\delta_g)^* = e_\beta(g)\delta_e.$$

Moreover, for $x \in E^1$ and $b \in e(x^*)\mathcal{B}$, we have

$$\varphi(\alpha_x(b)) = \varphi(\pi(x)b\pi(x)^*) = \rho(x)\varphi(b)\rho(x)^* = \beta_x(\varphi(b))\,,$$

showing that the restriction of φ to $\mathcal{B}$ is an equivariant homomorphism from $\mathcal{B}$ to $\mathcal{C}$. Observe that, being π a semi-saturated representation of $\mathbb{F}$ on $\mathcal{A}$, we have that $\alpha^\pi(g_1g_2\cdots g_n) = \alpha^\pi(g_1)\cdot\alpha^\pi(g_2)\cdot\cdots\cdot\alpha^\pi(g_n)$ in $\mathcal{I}(\mathbf{B})$, the inverse semigroup associated to the Boolean algebra $\mathbf{B}$ of $\mathcal{A}$, where $g_1g_2\cdots g_n$ is the reduced form of an element in $\mathbb{F}$. Since the corresponding statement for the partial representation ρ on $\mathcal{C} \rtimes_\beta \mathbb{F}$ also holds, we see that $\varphi(e(g)) = e_\beta(g)$ for all $g \in \mathbb{F}$. Now $\mathcal{B}$ is generated as an algebra by the projections $e(g)$, with $g \in \mathbb{F}$, and so we deduce from the above that $\varphi|_{\mathcal{B}}$ is the unique equivariant homomorphism from $\mathcal{B}$ to $\mathcal{C}$. We have shown that $(\mathcal{B}, \alpha)$ is a universal (E, C)-partial action.

Suppose conversely that $(\mathcal{B}, \alpha)$ is a universal partial (E, C)-action on the unital commutative $*$-algebra $\mathcal{B}$. Let α denote also the extension to a partial action of $\mathbb{F}$ on $\mathcal{B}$ described in Lemma 6.6, and let $\pi\colon \mathbb{F} \to \mathcal{B} \rtimes_\alpha \mathbb{F}$ be the corresponding semi-saturated partial representation (Lemma 6.6). Obviously, the unitaries $\pi(x)$, $x \in E^1$ satisfy (PI1)-(PI4). Let ρ be a semi-saturated partial representation of $\mathbb{F}$ on a unital $*$-algebra $\mathcal{A}'$ such that the partial isometries $\rho(x)$, $x \in E^1$, satisfy (PI1)-(PI4). Let $\mathcal{B}'$ be the subalgebra of $\mathcal{A}'$ generated by the projections $\rho(g)\rho(g)^*$, for $g \in \mathbb{F}$ and let β be the canonical partial action of $\mathbb{F}$ on $\mathcal{B}'$. Observe that, in particular, the partial action β induces a partial (E, C)-action on $\mathcal{B}'$. Therefore there is a unique equivariant homomorphism $\psi\colon \mathcal{B} \to \mathcal{B}'$. Since both π and ρ are semi-saturated we see that $\psi(e_\alpha(g)) = e_\beta(g)$ for all $g \in \mathbb{F}$, and that $\psi(\alpha_g(b)) = \beta_g(\psi(b))$ for all $b \in e_\alpha(g^{-1})\mathcal{B}$. We can thus define a $*$-homomorphism $\varphi\colon \mathcal{B} \rtimes_\alpha \mathbb{F} \to \mathcal{B}' \rtimes_\beta \mathbb{F}$ such that $\varphi(b\delta_g) = \psi(b)\delta_g$ for all $b \in e_\alpha(g)\mathcal{B}$. Composing the map $\varphi\colon \mathcal{B} \rtimes_\alpha \mathbb{F} \to \mathcal{B}' \rtimes_\beta \mathbb{F}$ with the natural $*$-homomorphism $\mathcal{B}' \rtimes_\beta \mathbb{F} \to \mathcal{A}'$ sending $b\delta_g$ to $b\rho(g)$ ([20, Proposition 6.8]), we obtain a $*$-homomorphism $\varphi'\colon \mathcal{B} \rtimes_\alpha \mathbb{F} \to \mathcal{A}'$ such that $\varphi'(\pi(g)) = \rho(g)$ for all $g \in \mathbb{F}$. If φ'' is another such $*$-homomorphism, then it must agree with φ' on the commutative subalgebra $\mathcal{B}$, and also

$$\varphi''(e_\alpha(g)\delta_g) = \varphi''(\pi(g)) = \rho(g) = \varphi'(e_\alpha(g)\delta_g)\,,$$

so that $\varphi' = \varphi''$.

We have shown that the partial representation π of $\mathbb{F}$ on $\mathcal{B} \rtimes_\alpha \mathbb{F}$ is universal for partial representations satisfying (PI1)-(PI4). This concludes the proof of the theorem. □

Note that we have a canonical partial (E,C) action on the unital commutative $*$-algebra B_∞, given by the projections $e(x)=xx^*$, $e(x^*)=s(x)$, and by the isomorphisms $\alpha_x\colon e(x^*)B_\infty \to e(x)B_\infty$ given by conjugation by $\underline{x}$, for all $x\in E^1$.

The following result follows immediately from Proposition 6.3 and Theorem 6.7.

Corollary 6.8. *The canonical partial (E,C)-action α on the unital commutative $*$-algebra B_∞ is universal.*

Similarly, we obtain the corresponding result for the completion $\mathcal{B}_0$ of B_∞:

Corollary 6.9. *The canonical partial (E,C)-action α on the unital commutative C*-algebra $\mathcal{B}_0$ is universal for partial (E,C)-actions on commutative C*-algebras.*

We can now use duality to obtain the universal property of the topological space $\Omega(E,C)$. Recall that the symbol $\sqcup$ is used to indicate disjoint unions.

Definition 6.10. An (E,C)-*dynamical system* consists of a compact Hausdorff space Ω with a family of clopen subsets $\{\Omega_v\}_{v\in E^0}$ such that

$$\Omega=\bigsqcup_{v\in E^0}\Omega_v,$$

and, for each $v\in E^{0,0}$, a family of clopen subsets $\{H_x\}_{x\in r^{-1}(v)}$ of Ω_v, such that

$$\Omega_v=\bigsqcup_{x\in X}H_x \qquad \text{for all } X\in C_v,$$

together with a family of homeomorphisms

$$\theta_x\colon \Omega_{s(x)}\longrightarrow H_x$$

for all $x\in E^1$.

Given two (E,C)-dynamical systems (Ω,θ), (Ω',θ'), there is an obvious definition of equivariant map $f\colon(\Omega,\theta)\to(\Omega',\theta')$, namely $f\colon\Omega\to\Omega'$ is *equivariant* if $f(\Omega_w)\subseteq\Omega'_w$ for all $w\in E^0$, $f(H_x)\subseteq H'_x$ for all $x\in E^1$ and $f(\theta_x(y))=\theta'_x(f(y))$ for all $y\in\Omega_{s(x)}$.

We say that an (E,C)-dynamical system (Ω,θ) is *universal* in case there is a unique continuous equivariant map from every (E,C)-dynamical system to (Ω,θ).

For instance if $(E,C)=(E(m,n),C(m,n))$, then this is exactly the definition of an (m,n)-dynamical system, given in [5, Definition 3.1].

Using the (contravariant) equivalence between commutative unital C*-algebras and compact Hausdorff topological spaces we get immediately from Corollary 6.9:

Corollary 6.11. *The dynamical system $(\Omega(E,C),\alpha^*)$ is the universal (E,C)-dynamical system.*

We also obtain the structure of $L_K^{\mathrm{ab}}(E,C)$ and $\mathcal{O}(E,C)$ as a crossed product:

Corollary 6.12. (1) *For any field with involution K, we have*

$$L_K^{\mathrm{ab}}(E,C) \cong C_K(\Omega(E,C)) \rtimes_\alpha \mathbb{F},$$

where $C_K(\Omega(E,C))$ is the algebra of K-valued continuous functions on $\Omega(E,C)$, and K is given the discrete topology.

(2) *We have*

$$\mathcal{O}(E,C) \cong C(\Omega(E,C)) \rtimes_\alpha \mathbb{F},$$

where $C(\Omega(E,C))$ is the C^*-algebra of continuous functions on $\Omega(E,C)$ (here $\mathbb{C}$ has the usual topology), and $C(\Omega(E,C)) \rtimes_\alpha \mathbb{F}$ denotes the full C^*-algebra crossed product.

7. The type semigroup

Let G be a group acting on a set X. Tarski's Theorem can be expressed as a result on the type semigroup $S(X,G)$, see [54, Theorem 9.1 and Corollary 9.2]. As Tarski shows in [51, Theorem 16.12], his result holds also if we replace total actions with partial actions. If G acts by continuous transformations on a topological space X, and $\mathbb{D}$ is a G-invariant subalgebra of subsets of X, then we may consider the corresponding type semigroup $S(X,G,\mathbb{D})$ as in [34], [35], [50]. As we will notice in this section, we may as well consider this semigroup for invariant subalgebras under a partial G-action. An important case, considered in the above references for global actions, is the case where X is a zero-dimensional metrizable compact space and $\mathbb{D}$ is the subalgebra of $\mathcal{P}(X)$ consisting of all the clopen subsets of X. It is an open question in this case whether the analogue of Tarski's Theorem holds. The proof of Tarski's Theorem is based on two special properties of the type semigroup $S(X,G)$: the Schröder-Bernstein axiom (also named Cantor-Schröder-Bernstein property) $x \le y$ and $y \le x$ imply $x = y$, and the cancellation property $nx = ny \implies x = y$. The Schröder-Bernstein axiom is known to hold for countably complete subalgebras. However the cancellation law is known to fail even for complete subalgebras [53].

Here we will show that for global actions of free groups on zero-dimensional metrizable compact spaces, the type semigroup $S(X,G,\mathbb{K})$, where $\mathbb{K}$ denotes the subalgebra of all clopen subsets of X, do not satisfy in general any cancellation condition (see Theorem 7.11). In particular we show in Corollary 7.12 the existence, for each pair of positive numbers (m,n) such that $1 < m < n$, of a global action of a finitely generated free group $\mathbb{F}$ on a zero-dimensional metrizable compact space Ω, such that Ω contains a non-paradoxical but (n,m)-paradoxical clopen subset. This is the first example of the failure of Tarski's Theorem in this setting.

Our main construction gives in a natural way *partial* actions on zero-dimensional metrizable compact spaces. Using globalization results from [1] we get corresponding global actions, although on a locally compact, non-compact, Hausdorff space. Passing finally to the one-point compactifications of these spaces, we obtain the desired global actions on zero-dimensional metrizable compact spaces.

Definition 7.1. Let $(\{X_t\}_{t\in G},\{\theta_t\}_{t\in G})$ be a partial action of a group G on a set X. Let $\mathbb{D}$ be a G-invariant subalgebra of $\mathcal{P}(X)$, with $X_t\in\mathbb{D}$ for all $t\in G$. Let $S(X,G,\mathbb{D})$ be the set

$$\Big\{\bigcup_{i=1}^{n} A_i\times\{i\}\mid A_i\in\mathbb{D}, n\in\mathbb{N}\Big\}\Big/\sim_S,$$

where the equivalence relation $\sim_S$ is defined as follows: two sets $A=\bigcup_{i=1}^{n}A_i\times\{i\}$ and $B=\bigcup_{j=1}^{m}B_j\times\{j\}$ are equivalent, denoted by $A\sim_S B$, if there exist $l\in\mathbb{N}$, $C_k\in\mathbb{D}$, $t_k\in G$, and natural numbers n_k,m_k, $k=1,\dots,l$, such that $C_k\subseteq X_{t^{-1}}$, and

$$A=\bigsqcup_{k=1}^{l}C_k\times\{n_k\},\qquad B=\bigsqcup_{k=1}^{t}\theta_t(C_k)\times\{m_k\}.$$

The equivalence class containing A is denoted by $[A]$, and addition is defined by

$$\Big[\bigcup_{i=1}^{n}A_i\times\{i\}\Big]+\Big[\bigcup_{j=1}^{m}B_j\times\{j\}\Big]=\Big[\bigcup_{i=1}^{n}A_i\times\{i\}\cup\bigcup_{j=1}^{m}B_j\times\{n+j\}\Big].$$

Note that $S(X,G,\mathbb{D})$ is a conical refinement monoid, with $0=[\emptyset]$. When $\mathbb{D}=\mathcal{P}(X)$, $S(X,G,\mathbb{D})$ is denoted by $S(X,G)$.

A subset E in $\mathbb{D}$ is said to be *G-paradoxical* (with respect to $\mathbb{D}$) in case the element $[E]$ of $S(X,G,\mathbb{D})$ is properly infinite, that is, $2[E]\le[E]$. With these definitions we can state Tarski's Theorem ([54, Corollary 9.2], [51, Theorem 16.12]) as follows:

Theorem 7.2. *Let $(\{X_t\}_{t\in G},\{\theta_t\}_{t\in G})$ be a partial action of a group G on a set X, and let E be a subset of X. Then E is G-non-paradoxical if and only if there exists a monoid homomorphism $\mu\colon S(X,G)\to\mathbb{R}^+\cup\{\infty\}$ such that $\mu([E])=1$.*

The proof of Tarski's Theorem is based on the following properties of $S(X,G)$ (see [54, Theorem 3.5 and Theorem 8.7]):

(a) *Schröder-Bernstein*: Let $x,y\in S(X,G)$. If $x\le y$ and $y\le x$ then $x=y$.

(b) *n-cancellation*: Let $n\ge 1$ and $x,y\in S(X,G)$. If $nx=ny$ then $x=y$.

For any semigroup satisfying these two properties, the two conditions stated in Tarski's Theorem are equivalent, that is, if e is a nonzero element in an abelian monoid S satisfying (a) and (b) above, then $2e\nleq e$ in S if and only if there is a monoid homomorphism $\mu\colon S\to\mathbb{R}^+\cup\{\infty\}$ such that $\mu(e)=1$. This is a consequence of a purely semigroup theoretic result [54, Theorem 9.1]. As observed in [50, Theorem 5.4], the two conditions stated in Tarski's Theorem are also equivalent if the abelian semigroup S satisfies the following condition:

(c) *almost unperforation*: $(n+1)x\le ny\implies x\le y$ for every $n\ge 1$ and $x,y\in S$.

We are interested in the following concrete setting. Let X be a zero-dimensional metrizable compact Hausdorff space, and let $\mathbb{K}=\mathbb{K}(X)$ be the subalgebra of $\mathcal{P}(X)$ consisting of all the clopen subsets of X. Let $(\{X_t\}_{t\in G},\{\theta_t\}_{t\in G})$ be a partial action of a group G by continuous transformations on X such that $X_t \in \mathbb{K}$ for all $t \in G$. Observe that $\mathbb{K}$ is then automatically G-invariant. Then we want to know to what extent properties (a),(b) or (c) hold in $S(X, G, \mathbb{K})$. In particular we want to know whether Tarski's Theorem holds in this context.

We have the following general result in this setting:

Lemma 7.3. *Let X be a totally disconnected compact Hausdorff topological space, and let $\mathbb{K}$ be the subalgebra of $\mathcal{P}(X)$ consisting of the clopen subsets of X. Let $(\{X_t\}_{t\in G}, \{\theta_t\}_{t\in G})$ be a continuous partial action of a group G on X, such that $X_t \in \mathbb{K}$ for all $t \in G$. Let K be a field, and let $C_K(X)$ be the K-algebra of K-valued continuous functions on X, where K is given the discrete topology. Then there exists a canonical monoid homomorphism*

$$\Phi\colon S(X, G, \mathbb{K}) \longrightarrow \mathcal{V}(C_K(X) \rtimes_{\theta^*} G)$$

sending $\left[\bigcup_{i=1}^n A_i \times \{i\}\right]$ *to* $\sum_{i=1}^n [1_{A_i}\delta_e]$*, for* $A_i \in \mathbb{K}$*,* $i = 1, \dots, n$*.*

Proof. Since additivity is clear, we only have to prove that the map is compatible with the equivalence $\sim_S$ of Definition 7.1. For this it is enough to show that if $C \in \mathbb{K}$ and $C \subseteq X_{t^{-1}}$ for some $t \in G$, then the projections $1_C\delta_e$ and $1_{\theta_t(C)}\delta_e$ are equivalent. Set $x = 1_{\theta_t(C)}\delta_t$, and note that $x^* = 1_C\delta_{t^{-1}}$. Then one computes that $xx^* = 1_{\theta_t(C)}\delta_e$ and $x^*x = 1_C\delta_e$, showing the result. □

We do not know whether the map defined in the lemma is injective or surjective in general, but we can show it is bijective for the actions of $\mathbb{F}$ on the spaces $\Omega(E, C)$ introduced in Section 6.

Theorem 7.4. *Let (E, C) be a finite bipartite separated graph such that $s(E^1) = E^{0,1}$ and $r(E^1) = E^{0,0}$, and let $(\Omega(E, C), \alpha^*)$ be the universal (E, C)-dynamical system (see Corollary 6.11). Then the map*

$$\Phi\colon S(\Omega(E, C), \mathbb{F}, \mathbb{K}) \longrightarrow \mathcal{V}(C_K(\Omega(E, C)) \rtimes_\alpha \mathbb{F})$$

is an isomorphism for any field K.

Proof. By Corollary 6.12(1) and Corollary 5.8, we have

$$\mathcal{V}(C_K(\Omega(E, C)) \rtimes_\alpha \mathbb{F}) \cong \mathcal{V}(L^{\mathrm{ab}}(E, C)) \cong \varinjlim(M(E_n, C^n), \iota_n) \cong M(F_\infty, D^\infty).$$

Let Φ' be the composition of the map Φ with the isomorphism $\mathcal{V}(C_K(\Omega(E, C)) \rtimes_\alpha \mathbb{F}) \cong \varinjlim(M(E_n, C^n), \iota_n)$. It will suffice to show that Φ' is an isomorphism.

Observe that there is a basis of clopen subsets of $\Omega(E, C)$ which is in bijective correspondence with the vertices in F_∞. Indeed, if $v \in D_n$ is a vertex in the n-th level, then it represents a minimal projection in the commutative algebra B_n. The image of this projection through the canonical limit map $B_n \to B_\infty$ is a projection in B_∞, and so corresponds to a clopen subset of $\Omega(E, C)$. Each clopen subset of $\Omega(E, C)$ is a finite (disjoint) union of clopen subsets coming from the vertices of E_n for some n. (Note that $E_n^0 = D_n \sqcup D_{n+1}$.) The map Φ' sends the class $[A_v]$ of the clopen set corresponding to a vertex $v \in D_n \sqcup D_{n+1}$ to the class in the limit $\varinjlim(M(E_n, C^n), \iota_n)$ of the corresponding element a_v of $M(E_n, C^n)$.

Put $S = S(\Omega(E,C),\mathbb{F},\mathbb{K})$. We are going to build an inverse monoid homomorphism $\Psi\colon \varinjlim(M(E_n,C^n),\iota_n) \to S$. To this end we need to define compatible monoid homomorphisms $\Psi_n\colon M(E_n,C^n) \to S$. For $v \in E_n^0$ define $\Psi_n(a_v) = [A_v]$, where, as before, A_v is the clopen subset of $\Omega(E,C)$ corresponding to v. If $X \in C_v$, with $v \in E_n^{0,0}$, then we have to show that the relation $a_v = \sum_{z\in X} a_{s(z)}$ is preserved by Ψ_n. Now, it follows from either (5.8) or (5.9) that, for $z \in X$, there is an element $u_z \in E^1 \cup (E^1)^*$ such that the image of the projection zz^* in $L^{\mathrm{ab}}(E,C) = A_\infty$ is a projection in B_∞ which is of the form $\alpha_{u_z}(1_{A_{s(z)}})$, where the clopen set $A_{s(z)}$ is contained in the domain of $\alpha^*_{u_z}$. Since $v = \sum_{z\in X} zz^*$ in $L(E_n,C^n)$, we conclude that

$$[A_v] = \sum_{z\in X}[\alpha^*_{u_z}(A_{s(z)})] = \sum_{z\in X}[A_{s(z)}], \tag{7.1}$$

showing that Ψ_n preserves the relation $a_v = \sum_{z\in X} a_{s(z)}$. To show compatibility with respect to $\iota_n\colon M(E_n,C^n) \to M(E_{n+1},C^{n+1})$, observe that, for $v \in E_n^{0,1}$, we have

$$\Psi_{n+1}(\iota_n(a_v)) = \Psi_{n+1}(a_v) = [A_v] = \Psi_n(a_v).$$

If $v \in E_n^{0,0}$ then, by the proof of Lemma 4.5(a),(b), we have $\iota_n(a_v) = \sum_{z\in X} a_{s(z)}$, where X is any element in C_v, so that using (7.1), we get

$$\Psi_{n+1}(\iota_n(a_v)) = \Psi_{n+1}\Big(\sum_{z\in X} a_{s(z)}\Big) = \sum_{z\in X}[A_{s(z)}] = [A_v] = \Psi_n(a_v),$$

as desired.

The homomorphism $\Psi\colon \varinjlim(M(E_n,C^n),\iota_n) \to S$ just defined is clearly a two-sided inverse of Φ'. This shows the result. ☐

We can now show our first main result on the type semigroup.

Theorem 7.5. *Let M be a finitely generated abelian conical monoid. Then there exist a zero-dimensional metrizable compact Hausdorff space Ω and a partial action of a finitely generated free group $\mathbb{F}$ on Ω such that there is a map $\iota\colon M \to S(\Omega,\mathbb{F},\mathbb{K})$ which is a refinement of M. In particular, the map $\iota\colon M \to S(\Omega,\mathbb{F},\mathbb{K})$ is a unitary embedding.*

Proof. By Redei's Theorem ([31]), M is finitely presented. Let $\langle \mathcal{X} \mid \mathcal{R}\rangle$ be a finite presentation of M, as in Definition 4.2, and let (E,C) be the associated finite bipartite separated graph, so that $M \cong M(E,C)$ (see Definition 4.2).

Let $\mathbb{F}$ be the free group on E^1, and let $(\Omega(E,C),\alpha^*)$ be the universal (E,C)-dynamical system (see Corollary 6.11). By Theorem 7.4, we have that the canonical map

$$\Phi\colon S(\Omega(E,C),\mathbb{F},\mathbb{K}) \longrightarrow \mathcal{V}(C_K(\Omega(E,C)) \rtimes_\alpha \mathbb{F})$$

is an isomorphism, and by Corollary 6.12(1) we have

$$\mathcal{V}(C_K(\Omega(E,C)) \rtimes_\alpha \mathbb{F}) \cong \mathcal{V}(L^{\mathrm{ab}}(E,C)).$$

Since the natural map $M(E,C) \to \mathcal{V}(L^{\mathrm{ab}}(E,C))$ is a refinement of $M(E,C)$ (Corollary 5.8), and $M \cong M(E,C)$, we obtain the result. ☐

In particular, we obtain the following:

Corollary 7.6. *There exist a partial action of a finitely generated free group $\mathbb{F}$ on a zero-dimensional metrizable compact space Ω, and a non-$\mathbb{F}$-paradoxical (with respect to $\mathbb{K}$) clopen subset A of Ω such that $\mu(A) = \infty$ for every nonzero $\mathbb{F}$-invariant finitely additive measure $\mu\colon \mathbb{K} \to [0, \infty]$.*

Proof. Take positive integers m, n such that $1 < m < n$, and set $(E, C) = (E(m, n), C(m, n))$. Set $\Omega := \Omega(E, C)$. Then there is a canonical partial action of the free group $\mathbb{F}$ on $m + n$ generators on the zero-dimensional metrizable compact space Ω. Moreover, it follows from Theorem 7.5 that there is a unitary embedding

$$\iota\colon \langle a \mid ma = na\rangle \longrightarrow S(\Omega, \mathbb{F}, \mathbb{K}).$$

Let $A = A_w$ be the clopen subset corresponding to the vertex w of the graph E. Then $[A]$ is an order-unit in $S(\Omega, \mathbb{F}, \mathbb{K})$, and $m[A] = n[A]$. Since $m < n$ this implies that $\mu(A) = \infty$ for every nonzero finitely additive $\mathbb{F}$-invariant measure μ defined on $\mathbb{K}$. Since $\iota(a) = [A]$ and $2a \nleq a$ in $\langle a \mid ma = na\rangle$, we deduce that $2[A] \nleq [A]$ in $S(\Omega, \mathbb{F}, \mathbb{K})$ (use properties (3) and (1) in Definition 3.7). It follows that A is not $\mathbb{F}$-paradoxical (with respect to $\mathbb{K}$). $\square$

We proceed now to extend the above results to the setting of global actions. This is accomplished by the use of the globalization techniques of [1].

Recall from [1, Example 1.1], that if

$$\beta : G \times Y \to Y$$

is an action of a given group G on a set Y, and if $X \subseteq Y$ is any subset, one may always restrict β to X, regardless of whether or not X is β-invariant, obtaining a *partial action*, as follows: for each $g \in G$, set $X_g = X \cap \beta_g(X)$, and observe that

$$\beta_g(X_{g^{-1}}) = \beta_g\big(X \cap \beta_{g^{-1}}(X)\big) = \beta_g(X) \cap X = X_g.$$

We may then define

$$\alpha_g : X_{g^{-1}} \to X_g$$

to be the restriction of β_g to $X_{g^{-1}}$, and it may be easily proven that α is a partial action of G on X, henceforth referred to as the *restriction of β to X*.

Conversely, if we start with a partial action α of G on X, a global (as opposed to partial) action β on a set $Y \supseteq X$ is called a *globalization*, or an *enveloping action* for α [1, Section 1], if α is the restriction of β to X, and moreover

$$Y = \bigcup_{g \in G} \beta_g(X).$$

Working in the category of topological spaces, as opposed to the category of sets, we always assume that the transformations α_g and β_g are continuous and that X is open in Y, in which case we will clearly have that each X_g is open in X.

It is our next goal to study the relationship between the type semigroup of a partial action α and that of its globalization or, more generally, any global action extending α.

For this let us assume we are given a set Y equipped with a fixed nonempty collection $\mathbb{E}$ of subsets of Y, which is closed under finite intersections, finite unions and relative complements. In other words, $\mathbb{E}$ is a *ring* of subsets of Y. From now on we will refer to the members of $\mathbb{E}$ as *admissible* subsets.

Let us assume that we are also given a fixed admissible subset $X \subseteq Y$, and let

$$\mathbb{E}|_X = \{E \cap X : E \in \mathbb{E}\} = \{D \in \mathbb{E} : D \subseteq X\}.$$

It is readily verified that $\mathbb{E}|_X$ is a subalgebra of subsets of X.

Let us now consider a global action β of a given group G on Y, which we will suppose to be *admissible* in the sense that $\beta_g(E)$ is admissible for every admissible subset $E \subseteq Y$. We may then consider the corresponding type semigroup $S(Y, G, \mathbb{E})$.

Denoting by α the restriction of β to X, it is clear that each X_g is admissible (relative to $\mathbb{E}|_X$) and that α is an *admissible partial action* in the sense that, if D is an admissible subset of some $X_{g^{-1}}$, then $\alpha_g(D)$ is admissible.

Given any $D \in \mathbb{E}|_X$, we shall denote the class of D in $S(X, G, \mathbb{E}|_X)$ by $[D]_X$. Likewise, the class in $S(Y, G, \mathbb{E})$ of any $E \in \mathbb{E}$ will be denoted by $[E]_Y$.

Proposition 7.7. *The correspondence*

$$\phi : [D]_X \mapsto [D]_Y, \quad (D \in \mathbb{E}|_X),$$

gives a well defined injective map from $S(X, G, \mathbb{E}|_X)$ to $S(Y, G, \mathbb{E})$.

Proof. We leave the easy verification of the well definedness of ϕ for the reader. In order to show that ϕ is injective, assume that $D, D' \in \mathbb{E}|_X$ are such that

$$[D]_Y = [D']_Y.$$

Then there are sets $D_1, \ldots, D_n \in \mathbb{E}$, such that $D = \bigsqcup_{i=1}^n D_i$ and, for suitable group elements $g_1, \ldots, g_n$, one has that $D' = \bigsqcup_{i=1}^n \beta_{g_i}(D_i)$.

Since the D_i are subsets of D, and hence also of X, it is evident that $D_i \in \mathbb{E}|_X$. Moreover notice that

$$\beta_{g_i}(D_i) \subseteq D' \cap \beta_{g_i}(X) \subseteq X \cap \beta_{g_i}(X) = X_{g_i},$$

so $D_i \subseteq X_{g_i^{-1}}$, and it thus makes sense to speak of $\alpha_{g_i}(D_i)$. It is also clear that

$$D' = \bigsqcup_{i=1}^n \alpha_{g_i}(D_i),$$

so $[D]_X = [D']_X$. The general case, that is, when D and D' are of the form $\bigcup_{i=1}^n D_i \times \{i\}$, may be treated similarly. □

A case of interest occurs when Y is a Hausdorff topological space and

$$\mathbb{E} = \mathbb{K}(Y) := \{E \subseteq Y : E \text{ is compact and open}\}.$$

Given a compact-open subset $X \subseteq Y$, notice that

$$\mathbb{K}(Y)|_X = \mathbb{K}(X).$$

Proposition 7.8. *Suppose we are given a continuous global action* β *of a discrete group* G *on a Hausdorff space* Y. *Also let* $X \subseteq Y$ *be a compact-open subset such that*

$$Y = \bigcup_{g \in G} \beta_g(X).$$

Letting α *be the restriction of* β *to* X, *one has that the map*

$$\phi : S\big(X, G, \mathbb{K}(X)\big) \to S\big(Y, G, \mathbb{K}(Y)\big)$$

described above is bijective.

Proof. Given $E \in \mathbb{K}(Y)$, we may view $\{\beta_g(X)\}_{g \in G}$ as an open covering of E. Observing that E is compact, there are finitely many group elements $g_1, \ldots, g_n$, such that $E \subseteq \bigcup_{i=1}^n \beta_{g_i}(X)$. Setting

$$\begin{aligned} E_1 &= E \cap \beta_{g_1}(X), \\ E_2 &= \big(E \cap \beta_{g_2}(X)\big) \setminus E_1, \\ E_3 &= \big(E \cap \beta_{g_3}(X)\big) \setminus (E_1 \cup E_2), \\ &\ \vdots \\ E_n &= \big(E \cap \beta_{g_n}(X)\big) \setminus (E_1 \cup \ldots \cup E_{n-1}) \end{aligned}$$

one has that $E = \bigsqcup_{i=1}^n E_i$. Observing that each $E_i \subseteq \beta_{g_i}(X)$, we have that

$$D_i := \beta_{g_i^{-1}}(E_i) \subseteq X.$$

Moreover, $\phi\big([D_i]_X\big) = [D_i]_Y = [E_i]_Y$, so

$$\phi\Big(\sum_{i=1}^n [D_i]_X\Big) = \sum_{i=1}^n \phi\big([D_i]_X\big) = \sum_{i=1}^n [E_i]_Y = [E]_Y.$$

Since $S\big(Y, G, \mathbb{K}(Y)\big)$ is generated by the collection of all elements of the form $[E]_Y$, as above, we deduce that ϕ is surjective. The fact that ϕ is injective follows from Proposition 7.7. ☐

Corollary 7.9. *Let* X *be a compact Hausdorff topological space which is totally disconnected in the sense that its topology admits a basis of compact-open sets. Also let* α *be a continuous partial action of the group* G *on* X. *Then* α *admits a unique globalization* β *on a totally disconnected, locally compact Hausdorff space* Y, *and moreover* $S\big(X, G, \mathbb{K}(X)\big)$ *is isomorphic to* $S\big(Y, G, \mathbb{K}(Y)\big)$.

Proof. By [1, Theorem 1.1] we know that α admits a unique globalization β acting on a topological space Y. Using [1, Proposition 1.2] we may easily show that Y is Hausdorff. Since X is open in Y, and $Y = \bigcup_{g \in G} \beta_g(X)$, it is clear that Y is locally compact and totally disconnected.

Finally, observing that X is a compact-open subset of Y, we may use Proposition 7.8 to conclude that $S\big(X, G, \mathbb{K}(X)\big)$ and $S\big(Y, G, \mathbb{K}(Y)\big)$ are isomorphic. ☐

We can now extend Theorem 7.4 to the globalized actions. For a totally disconnected locally compact Hausdorff space Y and a field K, we denote by $C_{c,K}(Y)$ the algebra of continuous functions with compact support from Y to K, where K is given the discrete topology. If θ is a global action of G on Y we get as in Lemma 7.3 a monoid homomorphism $\Phi\colon S(Y,G,\mathbb{K}(Y))\to\mathcal{V}(C_{c,K}(Y)\rtimes_{\theta^*}G)$.

Theorem 7.10. *Let (E,C) be a finite bipartite separated graph such that $s(E^1)=E^{0,1}$ and $r(E^1)=E^{0,0}$, and let $(\Omega(E,C),\alpha^*)$ be the universal (E,C)-dynamical system. Let $\Omega^{\mathfrak{G}}(E,C)$ denote the totally disconnected, locally compact Hausdorff space obtained from the globalization of the action of $\mathbb{F}$ on $\Omega(E,C)$ (Corollary 7.9), and let β^* be the corresponding global action of $\mathbb{F}$ on it.*

Then the map

$$\Phi\colon S(\Omega^{\mathfrak{G}}(E,C),\mathbb{F},\mathbb{K}(\Omega^{\mathfrak{G}}(E,C)))\longrightarrow\mathcal{V}(C_{c,K}(\Omega^{\mathfrak{G}}(E,C))\rtimes_{\beta}\mathbb{F})$$

is an isomorphism for any field K.

Proof. Set $X=\Omega(E,C)$ and $Y=\Omega^{\mathfrak{G}}(E,C)$. Let $e=1_X\in C_{c,K}(Y)$. Then it is easily seen that e is a full projection in $C_{c,K}(Y)\rtimes_\beta\mathbb{F}$ such that

$$e(C_{c,K}(Y)\rtimes_\beta\mathbb{F})e\cong C_K(X)\rtimes_\alpha\mathbb{F}.$$

It follows from the first paragraph of the proof of [6, Lemma 7.3] that the map

$$\mathcal{V}(\iota)\colon\mathcal{V}(C_K(X)\rtimes_\alpha\mathbb{F})\to\mathcal{V}(C_{c,K}(Y)\rtimes_\beta\mathbb{F})$$

induced by the inclusion $\iota\colon C_K(X)\rtimes_\alpha\mathbb{F}\to C_{c,K}(Y)\rtimes_\beta\mathbb{F}$ is an isomorphism. (See also [32, Corollary 5.6] for a more general statement.)

We have a commutative diagram:

$$\begin{tikzcd}
S(X,\mathbb{F},\mathbb{K}(X)) \arrow[r, "\Phi_X"] \arrow[d, "\cong"'] & \mathcal{V}(C_K(X)\rtimes_\alpha\mathbb{F}) \arrow[d, "\mathcal{V}(\iota)"', "\cong"] \\
S(Y,\mathbb{F},\mathbb{K}(Y)) \arrow[r, "\Phi_Y"] & \mathcal{V}(C_{c,K}(Y)\rtimes_\beta\mathbb{F})
\end{tikzcd}$$

We observed before that $\mathcal{V}(\iota)$ is an isomorphism. The map Φ_X is an isomorphism by Theorem 7.4, and the left vertical map is an isomorphism by Proposition 7.8. Therefore we conclude that Φ_Y is also an isomorphism. □

We finally can reach our goal of presenting global actions on zero-dimensional metrizable compact spaces with arbitrary failure of cancellation laws.

Theorem 7.11. *Let M be a finitely generated abelian conical monoid. Then there exist a zero-dimensional metrizable compact space Z and a global action of a finitely generated free group $\mathbb{F}$ on Z such that there is an order embedding $\iota\colon M\to S(Z,\mathbb{F},\mathbb{K}(Z))$.*

Proof. Let (E,C) be the finite bipartite separated graph considered in the proof of Theorem 7.5, and let $\mathbb{F}$ be the free group on E^1. Let $\Omega^{\mathfrak{G},\star}(E,C)=\Omega^{\mathfrak{G}}(E,C)\cup\{\star\}$ be the one-point compactification of the locally compact Hausdorff space $\Omega^{\mathfrak{G}}(E,C)$, with the action β^*

extended to $\Omega^{\mathfrak{G},\star}(E,C)$ by $\beta^*_t(\star)=\star$ for all $t\in\mathbb{F}$. Set $X=\Omega(E,C)$, $Y=\Omega^{\mathfrak{G}}(E,C)$ and $Z=\Omega^{\mathfrak{G},\star}(E,C)$.

By the proof of Theorem 7.5, there is an order-embedding $M\hookrightarrow S(X,\mathbb{F},\mathbb{K}(X))$, and, by Corollary 7.9, Y is a totally disconnected locally compact Hausdorff space such that the natural map

$$S(X,\mathbb{F},\mathbb{K}(X))\to S(Y,\mathbb{F},\mathbb{K}(Y))$$

is an isomorphism. It is easily seen that the natural map $S(Y,\mathbb{F},\mathbb{K}(Y))\to S(Z,\mathbb{F},\mathbb{K}(Z))$ is an order-embedding. Combining all these facts we get that Z is a zero-dimensional metrizable compact space and there is an order-embedding $M\hookrightarrow S(Z,\mathbb{F},\mathbb{K}(Z))$. □

Corollary 7.12. *There exist a global action of a finitely generated free group $\mathbb{F}$ on a zero-dimensional metrizable compact space Z, and a non-$\mathbb{F}$-paradoxical (with respect to $\mathbb{K}$) clopen subset A of Z such that $\mu(A)=\infty$ for every finitely additive $\mathbb{F}$-invariant measure $\mu\colon\mathbb{K}\to[0,\infty]$ such that $\mu(A)>0$.*

Proof. Consider the space $X=\Omega(E,C)$ and the clopen subset A appearing in the proof of Corollary 7.6, and set $Z=\Omega^{\mathfrak{G},\star}(E,C)$ as in the proof of Theorem 7.5. Then A is a clopen subset of Z satisfying the desired conditions. □

8. DESCRIPTIONS OF THE SPACE $\Omega(E,C)$.

We follow with the general hypothesis of Sections 4–6, so that (E,C) is a finite bipartite separated graph with $s(E^1)=E^{0,1}$ and $r(E^1)=E^{0,0}$. In this section we will provide descriptions of the zero-dimensional metrizable compact space $\Omega(E,C)$ and the partial action of the free group on E^1, $\mathbb{F}$, on the space $\Omega(E,C)$. These descriptions generalize the descriptions obtained in [5] for the universal (m,n)-dynamical system.

We will use the notation established in Definition 6.10, so that $\Omega(E,C)=\bigsqcup_{v\in E^0}\Omega(E,C)_v$, $\Omega(E,C)_v=\bigsqcup_{x\in X}H_x$ for all $X\in C_v$ $(v\in E^{0,0})$, and $\theta_x\colon\Omega(E,C)_{s(x)}\to H_x$ are the structural clopen sets and homeomorphisms of the universal (E,C)-dynamical system.

The main point to understand the structure of the space $\Omega(E,C)$ is the following: Points in $\Omega(E,C)$ are completely determined by their dynamics. Namely, suppose that we have a point ξ in $\Omega(E,C)_w$, where $w\in E^{0,1}$. Then we can apply the maps θ_x to ξ, for all $x\in E^1$ such that $s(x)=w$. Given one such arrow x, the point $\theta_x(\xi)$ belongs to $H_x\subseteq\Omega(E,C)_{r(x)}$. For each $X\in C_{r(x)}$ with $x\notin X$, we have a decomposition $\Omega(E,C)_{r(x)}=\bigsqcup_{y\in X}H_y$, and the point $\theta_x(\xi)$ belongs to exactly one of these sets H_y, say $\theta_x(\xi)\in H_{y(\xi,x)}$ for a unique $y(\xi,x)\in X$. Then we can apply the map $\theta^{-1}_{y(\xi,x)}$ to $\theta_x(\xi)$ and obtain a new point in $\Omega(E,C)_{s(y(\xi,x))}$, and again we can re-start the process. The point ξ is determined by the information provided by this infinite process, and conversely any coherent set of data determines a point in the space $\Omega(E,C)$.

It is easy to see that Proposition 6.3 generalizes to $\mathcal{O}(E,C)$, meaning that the latter is the universal C*-algebra for semi-saturated partial representations of $\mathbb{F}$ satisfying relations (PI1)–(PI4).

We may thus apply [26, Theorem 4.4] to obtain a characterization of $\Omega(E,C)$, the spectrum of relations (PI1)–(PI4), as a subset of $2^{\mathbb{F}}$. The first step is to translate our relations in terms of functions on $2^{\mathbb{F}}$. Relative to (PI1) we have the functions:

$$f^1_{x,y}(\xi) = [x\in\xi][y\in\xi] - \delta_{x,y}[x\in\xi], \quad \forall \xi\in 2^{\mathbb{F}}, \tag{F1}$$

for $x,y\in X$, $X\in C$, where the brackets correspond to boolean value. Speaking of (PI2), we have functions of the form:

$$f^2_{x,y}(\xi) = [x^{-1}\in\xi] - [y^{-1}\in\xi], \tag{F2}$$

for all $x,y\in E^1$ such that $s(x)=s(y)$. As for (PI3), the functions we consider are:

$$f^3_{X,Y}(\xi) = \sum_{x\in X}[x\in\xi] - \sum_{y\in Y}[y\in\xi], \tag{F3}$$

for all $X,Y\in C_v$, $v\in E^{0,0}$. In order to account for (PI4), we must consider a single function, namely:

$$f^4(\xi) = -1 + \sum_{w\in E^{0,1}}[x_w^{-1}\in\xi] + \sum_{v\in E^{0,0}}\sum_{x\in X_v}[x\in\xi], \tag{F4}$$

where we choose some x_w in $s^{-1}(w)$, for each $w\in E^{0,1}$, and some X_v in C_v, for each $v\in E^{0,0}$.

Finally, since $\mathcal{O}(E,C)$ is the universal C*-algebra for a collection of partial representations which are *semi-saturated*, we must include the functions

$$f^5_{g,h}(\xi) = [gh\in\xi][g\in\xi] - [gh\in\xi],$$

for all $g,h\in\mathbb{F}$, such that $|gh|=|g|+|h|$. See [22, Proposition 5.4] for more details.

Considering the set $\mathcal{R}$ formed by all of the above functions, we have by [26, Proposition 4.1], that

$$\Omega(E,C) = \Omega_{\mathcal{R}} = \{\xi\in 2^{\mathbb{F}} : 1\in\xi,\ f(g^{-1}\xi)=0,\ \text{for all } f\in\mathcal{R},\ \text{and all } g\in\xi\}.$$

In order to give a more explicit description for this space, let us first introduce some terminology. Given ξ in $2^{\mathbb{F}}$, and given g in ξ, we will let ξ_g be the *local configuration of* ξ *at* g, meaning the set of all elements

$$x\in E^1\cup(E^1)^{-1}$$

(recall that $\mathbb{F}$ is the free group generated by E^1), such that $gx\in\xi$. Thus the set $g\xi_g$ consists precisely of the elements in ξ whose distance to g equals 1 relative to the Cayley graph metric.

One may then check that $\Omega(E,C)$ consist precisely of all $\xi\in 2^{\mathbb{F}}$ such that:

- (a) $1\in\xi$,
- (b) ξ is convex, [25, Definition 4.4] and [25, Proposition 4.5],
- (c) for every $g\in\xi$, there is some $v\in E^0$ such that:
  - (c.1) in case $v\in E^{0,1}$, the local configuration ξ_g consists of the elements of the form e^{-1}, for all $e\in E^1$ such that $s(e)=v$ (and nothing more),
  - (c.2) in case $v\in E^{0,0}$, the local configuration ξ_g consists of exactly one element of each group $X\in C_v$ (and nothing more).

Having completed the description of $\Omega(E,C)$, the partial action of $\mathbb{F}$ is now easy to describe: for each $g \in \mathbb{F}$ we put

$$\Omega_g = \big\{\xi \in \Omega(E,C) : g \in \xi\big\},$$

and we let

$$\theta_g : \Omega_{g^{-1}} \to \Omega_g,$$

be given by $\theta_g(\xi) = g\xi = \{gh : h \in \xi\}$.

We now start the description of the space $\Omega(E,C)$ in terms of certain "choice functions".

To avoid cumbersome considerations, we will from now on assume the following:

Hypothesis 8.1. (E,C) is a finite bipartite separated graphs such that $|C_v| \geq 2$ for all $v \in E^{0,0}$ and $s(E^1) = E^{0,1}$.

This is justified by Proposition 9.2.

Given $e \in E^1$, there is a unique $X \in C$ such that $e \in X$. This unique X will be denoted by X_e.

Let $v \in E^{0,0}$. For $X \in C_v$, define

$$Z_X = \prod_{Y \in C_v,\, Y \neq X} Y.$$

For each $Y_0 \in C_v$, $Y_0 \neq X$, we denote by π_{Y_0} the canonical projection map $Z_X \to Y_0$.

We now provide a description of the points belonging to $\Omega(E,C)_w$ for $w \in E^{0,1}$.

Definition 8.2. A *partial E-function* for (E,C) is a finite sequence

$$(\Omega_1, f_1), (\Omega_2, f_2), \dots, (\Omega_r, f_r),$$

where each Ω_i is a certain finite subset of $\mathbb{F}$, and each f_i is a function from Ω_i to $\bigsqcup_{X \in C} Z_X$ satisfying the following rules:

1. $\Omega_1 = s^{-1}(w)$ for some $w \in E^{0,1}$ and $f_1(e) \in Z_{X_e}$ for all $e \in s^{-1}(w)$.
2. For each $j = 2, \dots, r$, Ω_j is the set of all the elements of $\mathbb{F}$ of the form

   $$e_{2j-1}e_{2j-2}^{-1}e_{2j-3}\cdots e_4^{-1}e_3e_2^{-1}e_1$$

   where

   $$e_{2j-3}\cdots e_4^{-1}e_3e_2^{-1}e_1 \in \Omega_{j-1},$$

   $e_{2j-2} = (\pi_Y \circ f_{j-1})(e_{2j-3}\cdots e_4^{-1}e_3e_2^{-1}e_1)$ for some $Y \in C$ which is a factor of Z_X, where $f_{j-1}(e_{2j-3}\cdots e_4^{-1}e_3e_2^{-1}e_1) \in Z_X$, and where $e_{2j-1} \in E^1$ satisfies that $s(e_{2j-1}) = s(e_{2j-2})$ and $e_{2j-1} \neq e_{2j-2}$.
3. The function $f_j \colon \Omega_j \to \bigsqcup_{X \in C} Z_X$ satisfies

   $$f_j(e_{2j-1}e_{2j-2}^{-1}e_{2j-3}\cdots e_4^{-1}e_3e_2^{-1}e_1) \in Z_{X_{e_{2j-1}}}$$

   for all $e_{2j-1}e_{2j-2}^{-1}e_{2j-3}\cdots e_4^{-1}e_3e_2^{-1}e_1 \in \Omega_j$. Note that, by Hypothesis 8.1, there is always at least one such function f_j.

An E-function is an infinite sequence $(\Omega_1, f_1), (\Omega_2, f_2), \dots$ satisfying the above conditions for all $j = 1, 2, 3, \dots$.

Note that it might happen that for some E-function $(\Omega_1, f_1), (\Omega_2, f_2), \dots$ we have $\Omega_r = \emptyset$ for some $r \geq 2$. In this case the E-function will be of the form

$$(\Omega_1, f_1), (\Omega_2, f_2), \dots, (\Omega_{r-1}, f_{r-1}), (\emptyset, \emptyset), (\emptyset, \emptyset), \dots.$$

This case occurs for instance for the separated graph (E, C) with $E^{0,0} = \{v\}$, $E^{0,1} = \{w_1, w_2\}$, and $C = C_v = \{X, Y\}$ with $X = \{\alpha\}$, $Y = \{\beta\}$, $s(\alpha) = s(\beta) = v$ and $r(\alpha) = w_1$, $r(\beta) = w_2$.

Note that every partial E-function $(\Omega_1, f_1), \dots, (\Omega_r, f_r)$ can be extended to an E-function $(\Omega_1, f_1), \dots, (\Omega_r, f_r), (\Omega_{r+1}, f_{r+1}), \dots$, although possibly $\Omega_i = \emptyset$ for all sufficiently large i. Another degenerate case is the case where there is only one extension of $(\Omega_1, f_1), \dots, (\Omega_r, f_r)$ to an E-function $(\Omega_1, f_1), \dots, (\Omega_r, f_r), (\Omega_{r+1}, f_{r+1}), \dots$, but $\Omega_i \neq \emptyset$ for all i. Note that this happens for instance for the unique partial E-function (Ω_1, f_1) of the separated graph $(E(1,1), C(1,1))$.

Theorem 8.3. *Let $\Omega(E, C)$ be the universal space associated to the separated graph (E, C), and let w be a vertex in $E^{0,1}$. Then the points of $\Omega(E, C)_w$ are in one-to-one correspondence with the E-functions $(\Omega_1, f_1), (\Omega_2, f_2), \dots$ such that $\Omega_1 = s^{-1}(w)$. Moreover, given a point ξ in $\Omega(E, C)_w$, represented by the E-function $(\Omega_1, f_1), (\Omega_2, f_2), \dots$, the partial E-functions $(\Omega_1, f_1), (\Omega_2, f_2), \dots, (\Omega_r, f_r)$, $r \geq 1$ represent a fundamental system of clopen subsets of ξ in $\Omega(E, C)$. The clopen set represented by the partial E-function $(\Omega_1, f_1), (\Omega_2, f_2), \dots, (\Omega_r, f_r)$ corresponds to a vertex in F_∞ lying in the $2r + 1$ level D_{2r+1} of $F^0_\infty = \bigsqcup_{i=0}^\infty D_i$.*

Proof. One can show, using arguments similar to those in [5, proof of Theorem 4.1] that the set of E-functions $(\Omega_1, f_1), (\Omega_2), f_2), \dots$, such that $\Omega_1 = s^{-1}(w)$ for $w \in E^{0,1}$, is in bijective correspondence with the set of configurations $\xi \in \Omega_{\mathcal{R}}$ such that the local configuration xi_1 of ξ at 1 consists of elements of the form e^{-1} for all $e \in s^{-1}(w)$ (form (c.1)).

We now proceed to show the statement about the relation between partial E-functions and vertices in the graph F_∞. First, we show that the partial E-functions (Ω_1, f_1) such that $\Omega_1 = s^{-1}(w)$ (for $w \in D_1$), correspond to vertices v in D_3 such that $r_3(v) = w$, where $r_n \colon D_n \to D_{n-2}$ is the map defined just before Lemma 5.2. Suppose that $\Omega_1 = s^{-1}(w) = \{x_1, \dots, x_r\}$. For $i = 1, \dots, r$, consider

$$y_i := \alpha^{x_i}(f_1(x_i)) \in E^1_1. \tag{8.1}$$

This has sense because $f_1(x_i) \in \prod_{Y \in C_{r(x_i)}, Y \neq X_{x_i}} Y$. Note that there are exactly r distinct elements in C_w, namely $X(x_1), \dots, X(x_r)$, and that $y_i \in X(x_i)$ for all i. We can therefore build the elements

$$z_i = \alpha^{y_i}(y_1, \dots \widehat{y_i}, \dots, y_r) \in E^1_2$$

for $i = 1, \dots, r$, and the vertex $v := v(y_1, \dots, y_r) \in D_3$. Note that $s^{-1}(v) = \{z_1, \dots, z_r\}$, so there is a canonical bijection $x_i \leftrightarrow z_i$ between $s^{-1}(w)$ and $s^{-1}(v)$. The vertex v just constructed is the vertex associated to the partial E-function (Ω_1, f_1). Conversely, the vertex $v = v(y_1, \dots, y_r)$ determines $y_1, \dots, y_r$, which in turn determine the function f_1 by using (8.1).

We will show that a partial E-function

$$(\Omega_1, f_1), \dots, (\Omega_r, f_r),$$

with $r \geq 1$, for (E, C), such that $\Omega_1 = s^{-1}(w)$, corresponds to a partial E-function

$$(\Omega'_1, f'_1), \dots, (\Omega'_{r-1}, f'_{r-1})$$

for the separated graph (E_2, C^2), such that $\Omega'_1 = s^{-1}(v)$, where v is the vertex in $D_3 = E_2^{0,1}$ corresponding to (Ω_1, f_1) (under the correspondence defined above). By induction hypothesis, this determines (and is determined by) a vertex v' in $(F'_\infty)^{0,2r-1}$, where F'_∞ is the F-graph associated to (E_2, C^2) (see Construction 4.4). Since F'_∞ is the separated graph obtained from F_∞ by eliminating its two first levels, we see that the original partial E-function corresponds to the vertex v' in $F_\infty^{0,2r+1} = D_{2r+1}$.

So we only need to prove the desired correspondence between partial E-functions for (E, C) and (E_2, C^2). Observe that, given a vertex v' in D_3 such that $r_3(v') = v \in D_1$, there is a canonical bijective map between $s^{-1}(v)$ and $s^{-1}(v')$. We will denote this bijection by $e \leftrightarrow e'$, for $e \in s^{-1}(v)$. Now take e in $s^{-1}(v)$. Then the corresponding arrow e' has an end vertex $r(e')$ and we are going to check that $r_2(r(e')) = r(e)$.

We have $s(e') = v(y_1, \dots, y_l)$ and $e' = \alpha^{y_t}(y_1, \dots, \widehat{y}_t, \dots, y_l)$ for some t, where $s^{-1}(s(e)) = \{x_1, \dots, x_l\}$ and $x_t = e$. Now set

$$C_{r(x_t)} = C_{r(e)} = \{X_1, \dots, X_s\}$$

and write $x_t = \widetilde{x}_p \in X_p$ for a unique $p \in \{1, \dots, s\}$. Then there are $\widetilde{x}_i \in X_i$ for $i \neq p$ such that

$$y_t = \alpha^{\widetilde{x}_p}(\widetilde{x}_1, \dots, \widehat{\widetilde{x}_p}, \dots, \widetilde{x}_s).$$

Note that $e \in X_p$, so that $X_e = X_p$. On the other hand

$$r(e') = r(\alpha^{y_t}(y_1, \dots, \widehat{y}_t, \dots, y_l)) = s(y_t) = v(\widetilde{x}_1, \dots, \widetilde{x}_s),$$

so that $r_2(r(e')) = r(e)$. It follows that there is a canonical bijective correspondence between $C_{r(e)}$ and $C_{r(e')}$ (see Lemma 5.2 and Remark 5.3), and we will denote this correspondence by $X \leftrightarrow X'$ for $X \in C_{r(e)}$. Note that $X_{e'} = X'_e$.

Let $(\Omega_1, f_1), \dots, (\Omega_r, f_r)$ be a partial E-function for (E, C), where $r > 1$. Let $w \in E^{0,1}$ such that $\Omega_1 = s^{-1}(w)$, and let w' be the vertex in D_3 corresponding to (Ω_1, f_1). We first define (Ω'_1, f'_1). Let $\Omega'_1 = \{e' \mid e \in \Omega_1\}$. Then f'_1 is determined as follows. Given $e' \in \Omega'_1$, there is a unique $e \in \Omega_1 = s^{-1}(w)$ corresponding to it through the bijection between $s^{-1}(w')$ and $s^{-1}(w)$. Now we are going to define $f'_1(e') \in Z_{X_{e'}}$. Adopt the notation of the above paragraph, so that $e' = \alpha^{y_t}(y_1, \dots, \widehat{y}_t, \dots, y_l)$ for some t, where $s^{-1}(s(e)) = \{x_1, \dots, x_l\}$ and $x_t = e$. Now set $C_{r(x_t)} = C_{r(e)} = \{X_1, \dots, X_s\}$ and write $e = \widetilde{x}_p \in X_p$ for a unique $p \in \{1, \dots, s\}$. Then there are $\widetilde{x}_i \in X_i$ for $i \neq p$ such that

$$y_t = \alpha^{\widetilde{x}_p}(\widetilde{x}_1, \dots, \widehat{\widetilde{x}_p}, \dots, \widetilde{x}_s).$$

For $i \neq p$, we have $X'_i \in C_{r(e')}$, and $X'_i \neq X'_p = X'_e = X_{e'}$, and we aim to define $\pi_{X'_i}(f'_1(e'))$. First set

$$\widetilde{y}_i = \alpha^{\widetilde{x}_i}(\widetilde{x}_1, \dots, \widehat{\widetilde{x}_i}, \dots, \widetilde{x}_s).$$

Observe that $r(\widetilde{y}_i) = s(\widetilde{x}_i)$. Moreover, note that it follows from (8.1) that $\widetilde{x}_i = \pi_{X_i}(f_1(e))$. The elements in $C_{r(\widetilde{y}_i)}$ are in bijective correspondence with the elements in $s^{-1}(s(\widetilde{x}_i))$. Write $\mathcal{S} = s^{-1}(s(\widetilde{x}_i)) \setminus \{\widetilde{x}_i\}$. Observe that, since $\widetilde{x}_i = \pi_{X_i}(f_1(e))$, the element $x\widetilde{x}_i^{-1}e$ belongs to Ω_2 for all $x \in \mathcal{S}$. Define $\pi_{X'_i}(f'_1(e')) = \alpha^{\widetilde{y}_i}((\widetilde{y}_{i,x})_{x\in\mathcal{S}})$, where

$$\widetilde{y}_{i,x} = \alpha^x(f_2(x\widetilde{x}_i^{-1}e)).$$

Note that, for X_i as above, with $i \neq p$, we have

$$r_3(s(\pi_{X'_i}(f'_1(e')))) = s(\widetilde{x}_i) = s(\pi_{X_i}(f_1(e))),$$

so that given any element $e'_3(\pi_{X'_i}(f'_1(e')))^{-1}e'$ in Ω'_2 we can form a corresponding element $e_3(\pi_{X_i}(f_1(e)))^{-1}e$ in Ω_2.

Assume that for some $r > j > 1$, we have defined a partial E-function $(\Omega'_1, f'_1), \dots, (\Omega'_{j-1}, f'_{j-1})$ on (E_2, C^2) satisfying the following conditions:

(1) For all $1 \leq j_0 < j$ and all $e'_{2j_0+1}(e'_{2j_0})^{-1} \cdots e'_3(e'_2)^{-1}e'_1$ in Ω'_{j_0+1} we have $r_3(s(e'_{2j_0})) = s(e_{2j_0})$, where e_{2j_0} is inductively defined by $e_{2j_0} = \pi_Y(f_{j_0}(e_{2j_0-1}e_{2j_0-2}^{-1} \cdots e_3e_2^{-1}e_1))$ if $e'_{2j_0} = \pi_{Y'}(f'_{j_0}(e'_{2j_0-1}(e'_{2j_0-2})^{-1} \cdots e'_3(e'_2)^{-1}e'_1))$ for some $Y \in C_{r(e_{2j_0-1})} \setminus \{X_{e_{2j_0-1}}\}$.

    Therefore we have a well-defined element $e_{2j_0+1}(e_{2j_0})^{-1} \cdots e_3e_2^{-1}e_1$ in Ω_{j_0+1} for each element $e'_{2j_0+1}(e'_{2j_0})^{-1} \cdots e'_3(e'_2)^{-1}e'_1$ in Ω'_{j_0+1}.

(2) For all $1 \leq j_0 < j$, all $e'_{2j_0-1}(e'_{2j_0-2})^{-1} \cdots e'_3(e'_2)^{-1}e'_1 \in \Omega'_{j_0}$, all $X \in C_{r(e_{2j_0-1})} \setminus \{X_{e_{2j_0-1}}\}$, and all $x \in s^{-1}\Big(s\Big(\pi_X f_{j_0}(e_{2j_0-1}e_{2j_0-2}^{-1} \cdots e_3e_2^{-1}e_1)\Big)\Big) \setminus \Big\{\pi_X f_{j_0}(e_{2j_0-1}e_{2j_0-2}^{-1} \cdots e_3e_2^{-1}e_1)\Big\}$, we have

$$\begin{aligned}&\pi_x\Big(\pi_{X'} f'_{j_0}(e'_{2j_0-1}(e'_{2j_0-2})^{-1} \cdots e'_3(e'_2)^{-1}e'_1)\Big)\\ &\quad = \alpha^x\Big(f_{j_0+1}\Big(x(\pi_X f_{j_0}(e_{2j_0-1} \cdots e_2^{-1}e_1))^{-1}e_{2j_0-1} \cdots e_3e_2^{-1}e_1\Big)\Big),\end{aligned}$$

    where π_x denotes the projection $\pi_{X(x)}$, being $X(x)$ the element in $C_{s(x)}$ corresponding to x.

Note that indeed condition (2) implies condition (1), but we found useful to state condition (1). As we have seen before, both conditions hold when $j_0 = 1$.

To complete the induction step, we have to define the map f'_j. This is done essentially as in the case $j = 1$, only the notation is a little bit worse.

Let $e'_{2j-1}(e'_{2j-2})^{-1} \cdots (e'_2)^{-1}e'_1$ be an element in Ω'_j, and let $e_{2j-1}e_{2j-2}^{-1} \cdots e_2e_1$ be the corresponding element in Ω_j (see condition (1)). We want to define $f'_j(e'_{2j-1}(e'_{2j-2})^{-1} \cdots (e'_2)^{-1}e'_1) \in Z_{X'_{e_{2j-1}}}$.

We have $s(e'_{2j-1}) = v(y_1, \dots, y_l)$ and $e'_{2j-1} = \alpha^{y_t}(y_1, \dots, \widehat{y_t}, \dots, y_l)$ for some t, where $s^{-1}(s(e_{2j-1})) = \{x_1, \dots, x_l\}$ and $x_t = e_{2j-1}$. Now set

$$C_{r(x_t)} = C_{r(e_{2j-1})} = \{X_1, \dots, X_k\}$$

and write $x_t = \widetilde{x}_p \in X_p$ for a unique $p \in \{1, \dots, k\}$. Then there are $\widetilde{x}_i \in X_i$ for $i \neq p$ such that

$$y_t = \alpha^{\widetilde{x}_p}(\widetilde{x}_1, \dots, \widehat{\widetilde{x}_p}, \dots, \widetilde{x}_k).$$

We have to check that $\widetilde{x}_i = \pi_{X_i}(f_j(e_{2j-1} \cdots e_2^{-1} e_1))$ for $i \neq p$. For this we need condition (2). Indeed, observe that there is $s \neq t$ such that

$$e'_{2j-2} = \alpha^{y_s}(y_1, \dots, \widehat{y_s}, \dots, y_l)$$

Observe that

$$e'_{2j-2} = \pi_{X_{e'_{2j-2}}}(f'_{j-1}(e'_{2j-3} \cdots (e'_2)^{-1} e'_1)).$$

On the other hand

$$\pi_{x_t}(e'_{2j-2}) = \pi_{x_t}(\alpha^{y_s}(y_1, \dots, \widehat{y_s}, \dots, y_l)) = y_t,$$

so that, putting everything together and using (2), we get

$$\begin{aligned} y_t &= \pi_{x_t}\pi_{X_{e'_{2j-2}}}(f'_{j-1}(e'_{2j-3} \cdots (e'_2)^{-1} e'_1)) \\ &= \alpha^{x_t}(f_j(e_{2j-1} e_{2j-2}^{-1} e_{2j-3} \cdots e_2^{-1} e_1)) \end{aligned}$$

because $e_{2j-2} = \pi_{X_{e_{2j-2}}}(f_{j-1}(e_{2j-3} \cdots e_2^{-1} e_1))$ and $e_{2j-1} = x_t$. This shows that, for $i \neq p$, we have $\widetilde{x}_i = \pi_{X_i}(f_j(e_{2j-1} \cdots e_2^{-1} e_1))$, as desired.

For $i \neq p$, we have $X'_i \in C_{r(e'_{2j-1})}$, and $X'_i \neq X'_p = X_{e'_{2j-1}}$, and we aim to define $\pi_{X'_i}(f'_j(e'_{2j-1} \cdots (e'_2)^{-1} e'_1))$. First set

$$\widetilde{y}_i = \alpha^{\widetilde{x}_i}(\widetilde{x}_1, \dots, \widehat{\widetilde{x}_i}, \dots, \widetilde{x}_k).$$

Observe that $r(\widetilde{y}_i) = s(\widetilde{x}_i)$.

The elements in $C_{r(\widetilde{y}_i)}$ are in bijective correspondence with the elements in $s^{-1}(s(\widetilde{x}_i))$. Write $\mathcal{S} = s^{-1}(s(\widetilde{x}_i)) \setminus \{\widetilde{x}_i\}$. Observe that, since $\widetilde{x}_i = \pi_{X_i}(f_j(e_{2j-1} \cdots e_2^{-1} e_1))$, the element $x\widetilde{x}_i^{-1} e_{2j-1} \cdots e_2^{-1} e_1$ belongs to Ω_{j+1} for all $x \in \mathcal{S}$. Define

$$\pi_{X'_i}(f'_j(e'_{2j-1} \cdots (e'_2)^{-1} e'_1)) = \alpha^{\widetilde{y}_i}((\widetilde{y}_{i,x})_{x \in \mathcal{S}}),$$

where

$$\widetilde{y}_{i,x} = \alpha^{x}(f_{j+1}(x\widetilde{x}_i^{-1} e_{2j-1} \cdots e_2^{-1} e_1)).$$

Note that, for X_i as above, with $i \neq p$, we have

$$r_3(s(\pi_{X'_i}(f'_j(e'_{2j-1} \cdots (e'_2)^{-1} e'_1))) = s(\widetilde{x}_i) = s(\pi_{X_i}(f_j(e_{2j-1} \cdots e_2^{-1} e_1))),$$

so that given any element $e'_{2j+1}(e'_{2j})^{-1} \cdots (e'_2)^{-1} e'_1$ in Ω'_{j+1} we can form a corresponding element $e_{2j+1} e_{2j}^{-1} \cdots e_2^{-1} e_1$ in Ω_{j+1}. This shows (1) for j, while (2) for j follows from the construction.

It follows from (2) that the partial E-function $(\Omega_1, f_1), \dots, (\Omega_r, f_r)$ is completely determined by $(\Omega'_1, f'_1), \dots, (\Omega'_{r-1}, f'_{r-1})$. (Notice that knowledge of the latter provides knowledge of the initial vertex w' in $D_3 = E_2^{0,1}$.)

Now observe that the points in F_∞ are in bijective correspondence with a basis of clopen subsets for the topology of $\Omega(E,C)$. Given an E-function $(\Omega_1, f_1), (\Omega_2, f_2), \dots$ we have a descending chain of clopen subsets $U_1 \supseteq U_2 \supseteq \cdots$, where U_r is the clopen set corresponding to $(\Omega_1, f_1), \dots, (\Omega_r, f_r)$. The intersection $\bigcap_{i=1}^\infty U_i$ will be non-empty, and using that $\Omega(E,C)$ is Hausdorff, it is easy to check that it consists of a single point ξ in $\Omega(E,C)$. This completes the proof of the theorem. ☐

The action of $\mathbb{F}$ on (partial) E-functions is described as follows:

Lemma 8.4. *Let*

$$g = z_r^{-1} x_r z_{r-1}^{-1} x_{r-1} \cdots z_1^{-1} x_1$$

be a reduced word in $\mathbb{F}$, *where* $x_1, \dots, x_r, z_1, \dots, z_r \in E^1$. *Then* $Dom(\theta_g) = \emptyset$ *unless* $z_i \in Y_i$ *for some* $Y_i \in C_{r(x_i)}$ *with* $Y_i \neq X_{x_i}$ *for all* $i = 1, \dots, r$, *and* $s(x_{i+1}) = s(z_i)$ *for* $i = 1, \dots, r-1$. *Assume that the latter conditions hold. Then the domain of* θ_g *is precisely the set of all* E*-functions* $\mathfrak{f} = ((\Omega_1, f_1), (\Omega_2, f_2), \dots)$ *such that* $\Omega_1 = s^{-1}(x_1)$ *and*

$$\pi_{Y_i} f_i(x_i \cdots z_2^{-1} x_2 z_1^{-1} x_1) = z_i$$

for all $i = 1, \dots, r$, *and the range of* g *is the set of those* E*-functions* $\mathfrak{f}' = (f'_1, f'_2, \dots,)$ *such that* $\pi_{X_{x_{r-i+1}}} f'_i(z_{r-i+1} \cdots z_{r-1} x_r^{-1} z_r) = x_{r-i+1}$ *for all* $i = 1, \dots, r$. *Moreover for* $\mathfrak{f} \in Dom(\theta_g)$ *let* $\mathfrak{f}' = {}^g\mathfrak{f}$ *denote the image of* $\mathfrak{f}$ *under the action of* g. *Then* $\mathfrak{f}' = ((\Omega'_1, f'_1), (\Omega'_2, f'_2), \dots)$ *with*

$$f'_{r+t}(y_{2t-1} \cdots y_2^{-1} y_1 x_1^{-1} z_1 \cdots z_{r-1} x_r^{-1} z_r) = f_t(y_{2t-1} \cdots y_2^{-1} y_1)$$

if $y_{2t-1} \cdots y_2^{-1} y_1 \in \Omega_t$ *and* $y_1 \neq x_1$.

Moreover, for $i = 2, 3, \dots, r$ *and* $y_{2t-1} \cdots y_2^{-1} y_1 z_{i-1}^{-1} x_{i-1} \cdots z_1^{-1} x_1 \in \Omega_{t+i-1}$ *with* $y_1 \neq x_i$,

$$f'_{r-i+1+t}(y_{2t-1} \cdots y_2^{-1} y_1 x_i^{-1} z_i \cdots z_{r-1} x_r^{-1} z_r) = f_{i-1+t}(y_{2t-1} \cdots y_2^{-1} y_1 z_{i-1}^{-1} x_{i-1} \cdots z_1^{-1} x_1),$$

and for $y_{2t-1} \cdots y_2^{-1} y_1 z_r^{-1} x_r \cdots z_2^{-1} x_2 z_1^{-1} x_1 \in \Omega_{t+r}$ *we have*

$$f'_t(y_{2t-1} \cdots y_2^{-1} y_1) = f_{r+t}(y_{2t-1} \cdots y_2^{-1} y_1 z_r^{-1} x_r \cdots z_2^{-1} x_2 z_1^{-1} x_1)\,.$$

Proof. The result is clear once we realize that the action of g on a point ξ in $\Omega(E,C)_{s(x_1)}$ consists in moving ξ using the composition $\theta_{z_r}^{-1} \theta_{x_r} \cdots \theta_{z_1}^{-1} \theta_{x_1}$, in case this is possible. ☐

9. EXAMPLES

Our first example is of a generic type. At first sight, it looks restrictive to concentrate attention to bipartite separated graphs. However, the following result shows that (modulo Morita equivalence) *all* graph C*-algebras (or Leavitt path algebras) can be studied that way. For a general separated graph (E,C) we define $L^{\mathrm{ab}}(E,C)$ (respectively $\mathcal{O}(E,C)$) as the quotient of $L(E,C)$ (respectively $C^*(E,C)$) by the ideal (resp. closed ideal) generated by the commutators $[e(u), e(u')]$, for all $u, u' \in U$, where U is the multiplicative semigroup generated by $E^1 \sqcup (E^1)^*$, and $e(u) = uu^*$.

Proposition 9.1. *Let (E,C) be a separated graph. Then there exists a bipartite separated graph $(\widetilde{E},\widetilde{C})$ such that*

$$L_K(\widetilde{E},\widetilde{C})\cong M_2(L_K(E,C)),\qquad C^*(\widetilde{E},\widetilde{C})\cong M_2(C^*(E,C)).$$

Moreover we have

$$L_K^{\mathrm{ab}}(\widetilde{E},\widetilde{C})\cong M_2(L_K^{\mathrm{ab}}(E,C)),\qquad \mathcal{O}(\widetilde{E},\widetilde{C})\cong M_2(\mathcal{O}(E,C)).$$

Proof. Let V_0 and V_1 be two disjoint copies of E^0, and denote the canonical maps $E^0\to V_i$ by $v\mapsto v_i$ for $i=0,1$. Write $\widetilde{E}^{0,0}=V_0$ and $\widetilde{E}^{0,1}=V_1$. Now $\widetilde{E}^1$ will be the disjoint union of a copy of E^0 and a copy of E^1:

$$\widetilde{E}^1=\{h_v\mid v\in E^0\}\bigsqcup\{e_0\mid e\in E^1\},$$

with

$$\tilde{r}(h_v)=v_0,\quad \tilde{s}(h_v)=v_1,\quad \tilde{r}(e_0)=r(e)_0,\quad \tilde{s}(e_0)=s(e)_1,\qquad (v\in E^0,e\in E^1).$$

Denote by e_{ij}, $0\le i,j\le 1$ the standard matrix units of $M_2(K)$. We define maps $\varphi\colon L(\widetilde{E},\widetilde{C})\to M_2(L(E,C))$ and $\psi\colon M_2(L(E,C))\to L(\widetilde{E},\widetilde{C})$ by the rules

$$\varphi(v_i)=v\otimes e_{ii},\quad \varphi(h_v)=v\otimes e_{01},\quad \varphi(e_0)=e\otimes e_{01},\qquad (v\in E^0,e\in E^1,i=0,1),$$

and

$$\begin{aligned}\psi(v\otimes e_{ii})&=v_i, &\psi(v\otimes e_{01})&=h_v, &&(v\in E^0,i=0,1),\\ \psi(e\otimes e_{00})&=e_0h^*_{s(e)}, &\psi(e\otimes e_{11})&=h^*_{r(e)}e_0 &&\\ \psi(e\otimes e_{01})&=e_0, &\psi(e\otimes e_{10})&=h^*_{r(e)}e_0h^*_{s(e)}, &&(e\in E^1).\end{aligned}$$

It is straightforward (though somewhat tedious) to check that φ and ψ give well-defined $*$-homomorphisms which are mutually inverse. Let U and $\widetilde{U}$ denote the multiplicative semigroups of the algebras $L(E,C)$ and $L(\widetilde{E},\widetilde{C})$ generated by $E^1\sqcup(E^1)^*$ and $\widetilde{E}^1\sqcup(\widetilde{E}^1)^*$ respectively. Let J denote the ideal of $L(E,C)$ generated by all elements of the form $[e(u),e(u')]$, with $u,u'\in U$, and let $\widetilde{J}$ denote the corresponding ideal of $L(\widetilde{E},\widetilde{C})$. Then it is easy to check that, for $\tilde{u}\in\widetilde{U}$, $\varphi(e(\tilde{u}))$ is a matrix consisting of an element of the form $e(u)$, for some $u\in U$, in one of the two main diagonal positions and 0s in the rest of positions. This clearly implies that $\varphi(\widetilde{J})\subseteq M_2(J)$. Since the image by ψ of an element of the form $u\otimes e_{00}$, for $u\in U$, belongs clearly to $\widetilde{U}$, we see that the reverse containment $M_2(J)\subseteq\varphi(\widetilde{J})$ also holds. Thus $\varphi(\widetilde{J})=M_2(J)$ and we conclude that φ defines an isomorphism from $L^{\mathrm{ab}}(\widetilde{E},\widetilde{C})$ onto $M_2(L^{\mathrm{ab}}(E,C))$, as desired.

The same proof applies (thanks to the universal properties of the involved C*-algebras) to show the corresponding result for the graph C*-algebras. □

In particular, since any classical graph C*-algebra $C^*(E)$ satisfies that the final projections $[e(u),e(u')]=0$ for all $u,u'\in U$, we see that $\mathcal{O}(E)=C^*(E)$ can always be interpreted (through a very concrete Morita equivalence) as a graph C*-algebra of a bipartite separated graph. A similar statement applies to Leavitt path algebras. The description of a Leavitt path algebra in terms of a crossed product by a partial action has been recently achieved by Gonçalves and Royer in [29].

We are now going to show that we can always reduce our study of the graph algebras of finite bipartite separated graphs, up to Morita equivalence and finite-dimensional summands, to the study of finite bipartite separated graphs satisfying Hypothesis 8.1.

Proposition 9.2. *Let (E, C) be a finite bipartite separated graph. Then there exists a finite bipartite separated graph (E', C') satisfying Hypothesis 8.1 such that $L_K(E, C)$ is Morita equivalent to $L_K(E', C') \oplus K^n$ for some $n \geq 0$. Similar statements hold for $C^*(E, C)$, $L^{\mathrm{ab}}(E, C)$ and $\mathcal{O}(E, C)$.*

Proof. Suppose that there is $v \in E^{0,0}$ such that $|C_v| = 1$. Let (F, D) be the bipartite separated graph such that $F^{0,0} = E^{0,0} \setminus \{v\}$, $F^{0,1} = E^{0,1}$, and $F^1 = E^1 \setminus r^{-1}(v)$. Set $p = \sum_{w \in E^0 \setminus \{v\}} w$. Then p is a full projection in $L_K(E, C)$ such that

$$L_K(F, D) \cong pL_K(E, C)p.$$

Therefore $L_K(E, C)$ is Morita equivalent to $L_K(F, D)$. We can apply this process to suppress all the vertices v' in $E^{0,0}$ such that $|C_{v'}| = 1$, and thus we arrive at a bipartite separated graph $(\widetilde{E}, \widetilde{C})$ such that $L_K(E, C)$ is Morita equivalent to $L_K(\widetilde{E}, \widetilde{C})$ and $|C_w| \neq 1$ for all $w \in E^{0,0}$.

Now let (E', C') be the bipartite separated graph obtained from $(\widetilde{E}, \widetilde{C})$ by deleting all the vertices in $\widetilde{E}^{0,0} \setminus r(\widetilde{E}^1)$ and all the vertices in $\widetilde{E}^{0,1} \setminus s(\widetilde{E}^1)$. Then clearly (E', C') satisfies Hypothesis 8.1, and $L_K(\widetilde{E}, \widetilde{C}) \cong L_K(E', C') \oplus K^n$, where $n = |E^0| - |r(\widetilde{E}^1)| - |s(\widetilde{E}^1)|$. This shows the result. ☐

We start our description of concrete examples with a motivational example for the entire theory of separated graphs, see [8], [7], [4], [5].

Example 9.3. For integers $1 \leq m \leq n$, define the separated graph $(E(m, n), C(m, n))$, where

(1) $E(m, n)^0 := \{v, w\}$ (with $v \neq w$).
(2) $E(m, n)^1 := \{\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_m\}$ (with $n + m$ distinct edges).
(3) $s(\alpha_i) = s(\beta_j) = v$ and $r(\alpha_i) = r(\beta_j) = w$ for all i, j.
(4) $C(m, n) = C(m, n)_v := \{X, Y\}$, where $X := \{\alpha_1, \dots, \alpha_n\}$ and $Y := \{\beta_1, \dots, \beta_m\}$.

See Figure 1 for a picture in the case $m = 2$, $n = 3$. By [8, Proposition 2.12],

$$L(E(m, n), C(m, n)) \cong M_{n+1}(L(m, n)) \cong M_{m+1}(L(m, n)),$$

where $L(m, n)$ is the classical Leavitt algebra of type (m, n). The same argument (by way of universal properties) shows that

$$C^*(E(m, n), C(m, n)) \cong M_{n+1}(U^{\mathrm{nc}}_{m,n}) \cong M_{m+1}(U^{\mathrm{nc}}_{m,n}),$$

where $U^{\mathrm{nc}}_{m,n}$ denotes the C*-algebra generated by the entries of a universal unitary $m \times n$ matrix, as studied by Brown and McClanahan in [15, 40, 41, 42].

The C*-algebras $\mathcal{O}_{m,n}$ and $\mathcal{O}^r_{m,n}$ studied in [5] are precisely the C*-algebras $\mathcal{O}(E(m, n), C(m, n))$ and $\mathcal{O}^r(E(m, n), C(m, n))$ in the notation of the present paper. (See Definition 10.7 for the definition of the reduced C*-algebra $\mathcal{O}^r(E,C)$.) It was shown in [5] that these two C*-algebras are not isomorphic.

The algebras $L^{\mathrm{ab}}_{m,n}(K) := L^{\mathrm{ab}}_K(E(m,n),C(m,n))$ have not appeared before in the literature. They provide natural examples (indeed universal examples) of actions on a compact Hausdorff space supporting (m,n)-paradoxical decompositions (Corollary 7.6).



Figure 1. The separated graph $(E(2,3),C(2,3))$

We now proceed to describe several concrete examples of separated graphs and their associated algebras. Curiously enough, these algebras (or the dynamical systems underlying them) have appeared before in the literature in different contexts. In many cases, one has to look at an appropriate full corner of the algebras to find the significant construction.

Example 9.4. Our first example is the "pure refinement" example, given in Figure 2. Let (E,C) be the separated graph described by that picture, with $C_v = \{X,Y\}$ and $X = \{\alpha_1,\alpha_2\}$, $Y = \{\beta_1,\beta_2\}$. The corner $vL_K(E,C)v$ is isomorphic to a free product $K^2 *_K K^2$. The corner of the abelianized Leavitt algebra $L^{\mathrm{ab}}_K(E,C)$ is isomorphic to K^4. We thus see a drastic reduction of the complexity in the transition from $L(E,C)$ to $L^{\mathrm{ab}}(E,C)$. A similar statement holds for the C*-algebras $C^*(E,C)$ and $\mathcal{O}(E,C)$.



Figure 2. The separated graph of pure refinement

Example 9.5. Let (E,C) be the separated graph described in Figure 3, with $C_v = \{X,Y\}$ and $X = \{\alpha_0,\alpha_1\}$ and $Y = \{\beta_0,\beta_1\}$.



Figure 3. The separated graph corresponding to Truss example

The Leavitt path algebra $L_K(E,C)$ is Morita equivalent to the corner $vL(E,C)v$, which is isomorphic to $M_2(K) *_K M_2(K)$.

By Corollary 6.12, $L_K^{\mathrm{ab}}(E,C) \cong C_K(\Omega(E,C)) \rtimes \mathbb{F}_4$, where $\Omega := \Omega(E,C)$ is a 0-dimensional compact space admitting a decomposition $\Omega = X \sqcup Y \sqcup Z$ into clopen subsets, such that Z decomposes in two different ways as a disjoint union of clopen subsets $Z = H_0 \sqcup H_1 = V_0 \sqcup V_1$, together with homeomorphisms $\mathfrak{h}_i \colon X \to H_i$ for $i = 0,1$, and $\mathfrak{v}_j \colon Y \to V_j$ for $j = 0,1$. This construction is closely related to Truss example in [53], which we briefly describe now. Let $\mathcal{E}$ be the non-directed graph with vertices $\{v_i \mid i \in \mathbb{Z}\}$ and with an edge connecting a vertex v_i with v_{i+1}, for all $i \in \mathbb{Z}$. Let T be the set of all the "labelings" of the edges and vertices of $\mathcal{E}$ satisfying the following conditions:

(i) the vertices are labelled 0 or 1, and adjacent vertices have different labels,
(ii) the edges are labelled by pairs (i,j) where $i,j \in \{0,1\}$,
(iii) if $(i,j),(k,l)$ are the labels on the edges incident with a vertex labelled 0 then $i \neq k$.
(iv) if $(i,j),(k,l)$ are the labels on the edges incident with a vertex labelled 1 then $j \neq l$.

Let

$$X = \{t \in T \mid v_0 \text{ is labelled } 0 \text{ and } (v_0,v_1) \text{ is labelled } (0,*)\}.$$

$$Y = \{t \in T \mid v_0 \text{ is labelled } 1 \text{ and } (v_0,v_1) \text{ is labelled } (*,1)\}.$$

Observe that (v_{-1},v_0) is labelled $(1,*)$ (respectively $(*,0)$) for every t in X (respectively Y).

Let T' be a disjoint copy of T, with vertices v_i' and with the same conditions on the labels of vertices and edges. Let $\iota \colon T \to T'$ be the canonical map. Set

$$Z = \{t' \in T' \mid v_0' \text{ is labelled } 0\}.$$

Consider the shift map $S \colon T \to T$ defined by the following conditions: $S(t)$ is the labelled graph such that the label of $S(t)$ at v_i is the label of t at v_{i-1}, and the label of $S(t)$ at (v_i,v_{i+1}) is the same as the label of t at (v_{i-1},v_i). Let $S' = \iota \circ S \circ \iota^{-1}$ be the corresponding bijection of T'.

We also need the involution $\phi \colon T \to T$, which sends any $t \in T$ to the labelled graph $\phi(t)$ defined by the condition that the bijection $v_i \to v_{-i}$ on the set of vertices $\{v_i \mid i \in \mathbb{Z}\}$ is an isomorphism of labelled graphs between t and $\phi(t)$.

We first indicate the connection between the definition using E-functions (see Theorem 8.3) and the description above. Consider an E-function $(\Omega_1,f_1),(\Omega_2,f_2),\ldots,$ where $\Omega_1 =$

$s^{-1}(w_0) = \{\alpha_0, \alpha_1\}$. This E-function will correspond, upon identification of the labels (i,j) with the labels (α_i, β_j), to an element t of X, as follows. The function f_1 takes values on $\{\beta_0, \beta_1\}$ so we define the label at the edge (v_0, v_1) as $(\alpha_0, f_1(\alpha_0))$ and the label at the edge (v_{-1}, v_0) as $(\alpha_1, f_1(\alpha_1))$. Now the label at the edge (v_1, v_2) is defined by $(f_2(\overline{f_1(\alpha_0)}f_1(\alpha_0)^{-1}\alpha_0), \overline{f_1(\alpha_0)})$, where we set $\overline{\alpha_i} = \alpha_{1-i}$ and $\overline{\beta_j} = \beta_{1-j}$. Similarly, the label at the edge (v_{-2}, v_{-1}) is defined by $(f_2(\overline{f_1(\alpha_1)}f_1(\alpha_1)^{-1}\alpha_1), \overline{f_1(\alpha_1)})$. Continuing in this way, we define the labels at all the edges (v_i, v_{i+1}). The labels at vertices are defined so as to make t an element of X. It is easy to see that this procedure establish a bijection between the set of E-functions $(\Omega_1, f_1), (\Omega_2, f_2), \dots$ with $\Omega_1 = s^{-1}(w_0)$ and X. Similarly there is a bijection between the set of E-functions $(\Omega_1, f_1), (\Omega_2, f_2), \dots$ with $\Omega_1 = s^{-1}(w_1)$ and Y.

We are now in a position to describe the maps $\mathfrak{h}_i, \mathfrak{v}_j$ in our picture of Ω. Set

$$H_0 = \{t' \in Z \mid (v_0', v_1') \text{ is labelled } (0,*)\}, \quad H_1 = \{t' \in Z \mid (v_0', v_1') \text{ is labelled } (1,*)\},$$

$$V_0 = \{t' \in Z \mid (v_0', v_1') \text{ is labelled } (*,0)\}, \quad V_1 = \{t' \in Z \mid (v_0', v_1') \text{ is labelled } (*,1)\},$$

Then $Z = H_0 \sqcup H_1 = V_0 \sqcup V_1$. Define homeomorphisms $\mathfrak{h}_0 \colon X \to H_0$ and $\mathfrak{h}_1 \colon X \to H_1$ by

$$\mathfrak{h}_0 = \iota|_X, \qquad \text{and} \qquad \mathfrak{h}_1 = (\iota \circ \phi)|_X.$$

Define homeomorphisms $\mathfrak{v}_0 \colon Y \to V_0$ and $\mathfrak{v}_1 \colon Y \to V_1$ by

$$\mathfrak{v}_0 = (S' \circ \iota)|_Y, \qquad \text{and} \qquad \mathfrak{v}_1 = (S' \circ \iota \circ \phi)|_Y.$$

Observe that $2[X] = 2[Y]$ in $S(\Omega, \mathbb{F}_4, \mathbb{E})$. Truss showed in [53] that $[X] \neq [Y]$ even in the semigroup $S(\Omega, \mathbb{F}_4, \mathbb{B})$, where $\mathbb{B}$ denotes the subalgebra of $\mathcal{P}(\Omega)$ consisting of the Borel subsets of Ω. Since $M(E,C) = \langle a, b \mid 2a = 2b \rangle$, the proof of Theorem 7.5 gives that $[X] \neq [Y]$ in $S(\Omega, \mathbb{F}_4, \mathbb{E})$.

Example 9.6. Let (E,C) be the separated graph described in Figure 4, with $C_v = \{X, Y\}$ and $X = \{\alpha_1, \alpha_2\}$ and $Y = \{\beta_1, \beta_2\}$. By [4, Lemma 5.5(a)], the corner $vC^*(E,C)v$ is the unital universal C*-algebra generated by a partial isometry. See [14] for a recent study of the *non-unital* universal C*-algebra generated by a partial isometry (of which $vC^*(E,C)v$ is the unitization).



FIGURE 4. The separated graph of a partial isometry

The C*-algebra $v\mathcal{O}(E,C)v$ is the C*-algebra of the *free monogenic inverse semigroup*, which was studied, among other places, in [33]. The 0-dimensional compact space $\Omega(E,C)$ decomposes as $\Omega(E,C) = X_v \sqcup \bigsqcup_{i=1}^{3} Y_{w_i}$, and we have two decompositions $X_v = H_{\beta_1} \sqcup H_{\beta_2} = H_{\alpha_1} \sqcup H_{\alpha_2}$ in clopen sets, with a universal homeomorphism $\alpha := \theta_{\alpha_1} \circ \theta_{\beta_1}^{-1} \colon H_{\beta_1} \to H_{\alpha_1}$, in the sense that given any other homeomorphism $\beta\colon X_1' \to Z_1'$, where X_1' and Z_1' are clopen subsets of a compact Hausdorff space X', there exists a unique equivariant continuous map from X' to X_v (cf. Corollary 6.11). The picture of the space X_v, together with the action of α on it is shown in Figure 5. Observe that the sets H_{β_2} and H_{α_2} are precisely the sets of points at the bottom row and right-most column, respectively. The unique fixed point of α is sited at the top left corner.



FIGURE 5. The compact space X_v.

We have $v\mathcal{O}(E,C)v \cong C(X_v) \rtimes_{\alpha^*} \mathbb{Z}$, which gives a possibly new description of this C*-algebra.

Example 9.7. Let (F,D) be the separated graph described in Figure 6, with $C_v = \{X,Y\}$ and $X = \{\alpha_1,\alpha_2\}$ and $Y = \{\beta_1,\beta_2\}$. By [4, Lemma 5.5(b)], we have

$$vC^*(E,C)v \cong C^*((*_{\mathbb{Z}}\mathbb{Z}_2) \rtimes \mathbb{Z})$$

where $\mathbb{Z}$ acts on $*_{\mathbb{Z}}\mathbb{Z}_2$ by shifting the factors of the free product. Similar computations give that $vL_K(E,C)v$ is isomorphic to the group algebra $K[(*_{\mathbb{Z}}\mathbb{Z}_2) \rtimes \mathbb{Z}]$.

It is easy to show that

$$v\mathcal{O}(F,D)v \cong C^*(\mathbb{Z}_2 \wr \mathbb{Z}), \qquad vL_K^{\mathrm{ab}}(F,D)v \cong K[\mathbb{Z}_2 \wr \mathbb{Z}],$$

where $\mathbb{Z}_2 \wr Z$ is the wreath product $(\oplus_{\mathbb{Z}}\mathbb{Z}_2) \rtimes \mathbb{Z}$. This is a well-known group, called the lamplighter group. This group provided the first counter-example to the Strong Atiyah's Conjecture, see [19], [30]. Observe that $\Omega(E,C) = X_v \sqcup Y_{w_1} \sqcup Y_{w_2}$, with $X_v \cong \prod_{\mathbb{Z}} \mathbb{Z}_2$ being the *Pontrjagin dual* of $\oplus_{\mathbb{Z}}\mathbb{Z}_2$. Here we have two decompositions $X_v = H_{\beta_1} \sqcup H_{\beta_2} = H_{\alpha_1} \sqcup H_{\alpha_2}$



FIGURE 6. The separated graph underlying the lamplighter group

in clopen sets, and a universal homeomorphism α of X_v sending H_{β_i} to H_{α_i}, for $i = 1, 2$. The action of α on X_v is essentially the dual action of the action of $\mathbb{Z}$ on $\oplus_{\mathbb{Z}}\mathbb{Z}_2$. We obtain the well-known representation of the group algebra of the lamplighter group:

$$K[\mathbb{Z}_2 \wr \mathbb{Z}] \cong vL_K^{\mathrm{ab}}(F, D)v \cong C_K\Big(\prod_{\mathbb{Z}} \mathbb{Z}_2\Big) \rtimes \mathbb{Z}.$$

The group algebras $K[\mathbb{Z}_p \wr \mathbb{Z}]$, $p \geq 3$, can be similarly represented, using a bipartite separated graph with $p+1$ vertices $v, w_1, \dots, w_p$, with $E^{0,0} = \{v\}$, $E^{0,1} = \{w_1, \dots, w_p\}$, and $2p$ arrows $\{\alpha_i, \beta_i \mid i = 1, \dots, p\}$, with $s(\alpha_i) = s(\beta_i) = w_i$ and $C_v = \{X, Y\}$, for $X = \{\alpha_1, \dots, \alpha_p\}$, $Y = \{\beta_1, \dots, \beta_p\}$.

10. TOPOLOGICALLY FREE ACTIONS

In this final section we characterize the finite bipartite separated graphs (E, C) such that the canonical action of the free group $\mathbb{F}$ on the universal space $\Omega(E, C)$ is topologically free, and we obtain consequences for the algebraic structure of the reduced C*-algebra $\mathcal{O}^r(E, C)$ (see Definition 10.7) and of the abelianized Leavitt algebra $L^{\mathrm{ab}}(E, C)$.

Recall the following definition from [26].

Definition 10.1. Let θ be a partial action of a group G on a compact Hausdorff space X. The partial action θ is *topologically free* if for every $t \in G \setminus \{1\}$, the set $F_t := \{x \in U_{t^{-1}} \mid \theta_t(x) = x\}$ has empty interior.

We now give our particular version of condition (L).

We will use paths in the *double* of E, that is, the graph $\hat{E}$ obtained from E by adjoining, for each $e \in E^1$, an edge e^* going in the reverse direction of e, that is $s(e^*) = r(e)$ and $r(e^*) = s(e)$. Given a path

$$\gamma = e_{2r}^* e_{2r-1} \cdots e_4^* e_3 e_2^* e_1,$$

in $\hat{E}$, we say that γ is an *admissible path* in case $X_{e_{2i-1}} \neq X_{e_{2i}}$ for $i = 1, 2 \dots, r$, that is, the edges e_{2i-1} and e_{2i} belong to different elements of $C_{r(e_{2i-1})} = C_{r(e_{2i})}$ for all $i = 1, 2 \dots, r$, and $e_{2i+1} \neq e_{2i}$ for $i = 1, \dots, r-1$. The same definition applies to paths of the forms $e_{2r-1} \cdots e_4^* e_3 e_2^* e_1$, or $e_{2r}^* e_{2r-1} \cdots e_4^* e_3 e_2^*$, or $e_{2r-1} \cdots e_4^* e_3 e_2^*$.

Definition 10.2. Let (E, C) be a finite bipartite separated graph, with $s(E^1) = E^{0,1}$ and $r(E^1) = E^{0,0}$. A *closed path* in (E, C) is a non-trivial admissible path $\gamma = e_{2r}^* e_{2r-1} \cdots e_2^* e_1$ such that $s(e_1) = s(e_{2r})$.

A *cycle* in (E, C) is a closed path $\gamma = e_{2r}^* e_{2r-1} \cdots e_2^* e_1$ in (E, C) such that $e_1 \neq e_{2r}$ and $s(e_{2i-1}) \neq s(e_{2j})$ for all $1 \leq i \leq j < r$.

Let γ be a closed path in (E, C). An *entry* of γ is a non-trivial admissible path of the form $x_{2t-1} x_{2t-2}^* \cdots x_4^* x_3 x_2^* x_1$ such that $s(x_1) = s(e_1)$, and $|X| \geq 2$ for some $X \in C_{r(x_{2t-1})}$ such that $X \neq X_{x_{2t-1}}$.

We say that (E, C) *satisfies condition (L)* if every cycle has an entry.

Note that condition (L) is very common. Indeed, for the separated graphs (E, C) associated to one-relator monoids

$$\langle a_1, a_2, \dots, a_n \mid \sum_{i=1}^n r_i a_i = \sum_{i=1}^n s_i a_i \rangle$$

with r_i, s_i non-negative integers such that $r_i + s_i >$ for all i, $\sum_i r_i > 0$, and $\sum_i s_i > 0$ (see [4]), the only one that does not satisfy condition (L) is the one corresponding to the presentation $\langle a \mid a = a \rangle$. Its separated graph is the $(E(1,1), C(1,1))$. Obviously, the cycle $\beta^* \alpha$ does not have exits.

We need a couple of preparatory lemmas.

Lemma 10.3. *Let $\gamma = e_{2r}^* e_{2r-1} \cdots e_2^* e_1$ be a closed path in (E, C), and assume that, for some $1 \leq i < r$ there exists an admissible path $\nu = x_{2t-1} x_{2t}^* \cdots x_4^* x_3 x_2^* x_1$ such that $s(x_1) = s(e_{2i})$ and $|X| \geq 2$ for some $X \in C_{r(x_{2t-1})}$ such that $X \neq X_{x_{2t-1}}$. Then there exists an entry for γ.*

Proof. If $x_1 \neq e_{2i}$ then $\nu e_{2i}^* e_{2i-1} \cdots e_2^* e_1$ is an exit for γ.

If $t \geq i+1$ and $x_1 = e_{2i}, x_2 = e_{2i-1}, \dots, x_{2i} = e_1$, then $x_{2t-1} x_{2t-2}^* \cdots x_{2i+2}^* x_{2i+1}$ is an entry for γ.

Suppose that $x_1 = e_{2i}, x_2 = e_{2i-1}, \dots, x_{2k} = e_{2i-2k+1}$, but $x_{2k+1} \neq e_{2i-2k}$ for some $k < i$. Then

$$x_{2t-1} x_{2t-2}^* \cdots x_{2k+1} e_{2i-2k}^* e_{2i-2k-1} \cdots e_2^* e_1$$

is an exit for γ.

Suppose that $x_1 = e_{2i}, x_2 = e_{2i-1}, \dots, x_{2k} = e_{2i-2k+1}$, $x_{2k+1} = e_{2i-2k}$, for some $k < i$, and that either $2t-1 = 2k+1$ or $x_{2k+2} \neq e_{2i-2k-1}$.

If $2k+1 = 2t-1$ then

$$\nu = x_{2k+1} x_{2k}^* \cdots x_2^* x_1 = e_{2i-2k} e_{2i-2k+1}^* \cdots e_{2i-1}^* e_{2i}.$$

There exists $X \in C_{r(e_{2i-2k-1})}$ such that $|X| \geq 2$ and $e_{2i-2k} \notin X$. We distinguish two cases:

(a) If $e_{2i-2k-1} \notin X$, then $e_{2i-2k-1} e_{2i-2k-2}^* \cdots e_2^* e_1$ is an entry for γ.

(b) If $e_{2i-2k-1} \in X$, then $e_{2i-2k} e_{2i-2k+1}^* \cdots e_{2r-1}^* e_{2r}$ is an entry for γ.

Assume now that $x_{2k+2} \neq e_{2i-2k-1}$, and write

$$\nu = \nu' x_{2k+1} x_{2k}^* \cdots x_2^* x_1$$

with ν' of positive length. Again we distinguish two cases:

(a) x_{2k+2} and $e_{2i-2k-1}$ belong to different sets of $C_{r(e_{2i-2k-1})}$. In this case the admissible path

$$\nu' x_{2k+2}^* e_{2i-2k-1} e_{2i-2k-2}^* \cdots e_2^* e_1$$

is an entry for γ.

(b) x_{2k+2} and $e_{2i-2k-1}$ belong to $X \in C_{r(e_{2i-2k-1})}$. In particular, we have $|X| \geq 2$, because we are assuming that $x_{2k+2} \neq e_{2i-2k-1}$. It follows that

$$e_{2i-2k} e_{2i-2k+1}^* \cdots e_{2r-1}^* e_{2r}$$

is an entry for γ.

□

Lemma 10.4. *Let $\gamma = e_{2r}^* e_{2r-1} \cdots e_2^* e_1$ be a closed path in (E, C), and suppose that (E, C) satisfies condition (L). Then the path γ has an entry.*

Proof. Suppose that there are $1 \leq i \leq j < r$ such that $s(e_{2i-1}) = s(e_{2j})$. Taking such a pair (i, j) with $j - i$ minimal, we have that $s(e_{2k-1}) \neq s(e_{2l})$ for all k, l such that $i \leq k \leq l < j$. Using Lemma 10.3, and changing notation, we can thus assume that $s(e_{2i-1}) \neq s(e_{2j})$ for $1 \leq i \leq j < r$.

If in addition $e_1 \neq e_{2r}$, then γ is a cycle and so there is an entry for γ by condition (L).

Suppose that $e_1 = e_{2r}$. Note that this implies $r > 1$ because γ is an admissible path. We claim that $e_{2r-1} \neq e_2$. If $r = 2$ then $e_2 \neq e_3 = e_{2r-1}$. If $r \geq 3$, then

$$s(e_2) = s(e_3) \neq s(e_{2(r-1)}) = s(e_{2r-1})$$

and consequently $e_2 \neq e_{2r-1}$. Therefore $e_2 \neq e_{2r-1}$ in any case. Write $X = X_{e_{2r-1}}$, $Y = X_{e_1} = X_{e_{2r}}$, $Z = X_{e_2}$. Then $X \neq Y$ and $Y \neq Z$, and $X, Y, Z \in C_{r(e_1)}$. There are two cases: $X = Z$ and $X \neq Z$. If $X = Z$, then since $e_2, e_{2r-1} \in X$ and $e_2 \neq e_{2r-1}$ we have $|X| \geq 2$, and thus e_1 is an entry for γ. If $X \neq Z$, then

$$e_2^* e_{2r-1} e_{2r-2}^* e_{2r-3} \cdots e_6^* e_5 e_4^* e_3$$

is a cycle, and consequently it has an entry η, because (E, C) has condition (L). It follows from Lemma 10.3 that γ has an entry. □

Theorem 10.5. *Let (E, C) be a finite bipartite separated graph, with $s(E^1) = E^{0,1}$ and $r(E^1) = E^{0,0}$. Then the partial action of $\mathbb{F}$ on $\Omega(E, C)$ is topologically free if and only if (E, C) satisfies condition (L).*

Proof. Suppose first that (E, C) does not satisfy condition (L), and let

$$\gamma = e_{2r}^* e_{2r-1} \cdots e_2^* e_1$$

by a cycle without exits in (E, C). Write

$$g = e_{2r}^{-1} e_{2r-1} \cdots e_2^{-1} e_1 \in \mathbb{F}.$$

Note that $g \neq 1$, because the defining expression of g is the reduced form of g (with respect to the canonical generating set E^1). Observe that it follows from the condition that the cycle γ does not have exits that there is a unique E-function $(\Omega_1, f_1), (\Omega_2, f_2), \dots$ such that $\Omega_1 = s(e_1)$. Let ξ be the point of $\Omega(E, C)$ represented by this E-function (see Theorem 8.3). Then we have $\Omega(E, C)_{s(e_1)} = \{\xi\}$.

Since γ is a cycle, it follows that $\theta_g(\xi) = \xi$. Therefore $\Omega(E, C)_{s(e_1)} = \{\xi\}$ is a clopen subset of $\Omega(E, C)$, fixed by the non-trivial element g of $\mathbb{F}$.

Conversely, suppose that (E, C) satisfies condition (L). Assume first that

$$g = z_r^{-1} x_r z_{r-1}^{-1} x_{r-1} \cdots z_1^{-1} x_1$$

is a reduced word in $\mathbb{F} \setminus \{1\}$. We may assume that $\mathrm{Dom}(\theta_g) \neq \emptyset$, and this implies by Lemma 8.4 that

$$\gamma = z_r^* x_r z_{r-1}^* x_{r-1} \cdots z_1^* x_1$$

is an admissible path in (E, C). We may also assume that θ_g has some fixed point. So let ξ be a fixed point of θ_g, with associated E-function $\mathfrak{f} = (\Omega_1, f_1), (\Omega_2, f_2), \dots$. By Lemma 8.4, we have $\Omega_1 = s^{-1}(s(x_1))$. Let $\mathfrak{f}' = ((\Omega_1', f_1'), (\Omega_2', f_2'), \dots)$ be the E-function corresponding to $\theta_g(\xi)$. Since $\theta_g(\xi) = \xi$, we must have $\Omega_i' = \Omega_i$ and $f_i' = f_i$ for all i. In particular, since $\Omega_1' = s^{-1}(s(z_r))$, we must have $s(x_1) = s(z_r)$. It follows that γ is a closed path in (E, C).

Suppose first that $x_1 \neq z_r$. Note that then

$$\gamma^i = z_r^* x_r \cdots z_1^* x_1 z_r^* x_r \cdots z_2^* x_2 z_1^* x_1$$

is a closed path in (E, C). Let W be an open neighborhood of ξ. By Theorem 8.3, there exists a positive integer $k \geq r$ such that W contains the clopen neighborhood U of ξ defined by the partial E-function $(\Omega_1, f_1), \dots, (\Omega_k, f_k)$. We will show that U contains a point ξ' such that $\theta_g(\xi') \neq \xi'$. By Lemma 10.4, there exists an entry

$$\eta = y_{2t-1} y_{2t-2}^* y_{2t-1} \cdots y_2^* y_1$$

for γ. By definition, this means that η is an admissible path such that $s(y_1) = s(x_1)$ and such that there is $X \in C_{r(y_{2t-1})}$ such that $|X| \geq 2$ and $y_{2t-1} \notin X$. We may assume that t is minimal with this property, so that for every $1 \leq t' < t$, we have $|X'| = 1$ for all $X' \in C_{r(y_{2t'-1})}$ such that $y_{2t'-1} \notin X'$. Observe that this implies that $\eta \in \Omega_t$, because for every $X' \in C_{r(y_{2t'-1})}$ such that $X_{y_{2t'-1}} \neq X'$ we have that $\pi_{X'} f_{t'}(y_{2t'-1} y_{2t'-2}^{-1} \cdots y_2^{-1} y_1)$ is the unique element of X', and so in particular we have that

$$y_{2t'} = \pi_{X_{y_{2t'}}} f_{t'}(y_{2t'-1} y_{2t'-2}^{-1} \cdots y_2^{-1} y_1).$$

Write $g_1 = y_{2t-1} y_{2t-2}^{-1} \cdots y_2^{-1} y_1 \in \mathbb{F}$.

Now let i be a positive integer such that $ir + t > k$. If $x_1 \neq y_1$, then

$$\eta(\gamma^*)^i = y_{2t-1} y_{2t-2}^* \cdots y_2^* y_1 x_1^* z_1 \cdots x_r^* z_r x_1^* z_1 \cdots \quad \cdots x_r^* z_r$$

is an admissible path. Now observe that, by Lemma 8.4, we have $g_1 g^{-i} \in \Omega_{ir+t}' = \Omega_{ir+t}$ and

$$f_{ir+t}(g_1 g^{-i}) = f_{ir+t}'(g_1 g^{-i}) = f_t(g_1).$$

Now let ξ' be the E-function $(\Omega''_1, f''_1), (\Omega''_2, f''_2), \ldots$ defined as follows. For $1 \leq j \leq ir + t - 1$, set $\Omega''_j = \Omega_j$ and $f''_j = f_j$. Now let $X \in C_{r(y_{2t-1})}$ such that $|X| \geq 2$ and $y_{2t-1} \notin X$. This means that there is $x \in X$ such that

$$x \neq f_t(y_{2t-1}y_{2t-2}^{-1}\cdots y_2^{-1}y_1) = f_t(g_1).$$

We define

$$\pi_X f''_{ir+t}(g_1 g^{-i}) = x,$$

and we complete the definition of f''_{ir+t} arbitrarily, so that we get a partial E-function $(\Omega''_1, f''_1), \ldots (\Omega''_{ir+t}, f''_{ir+t})$. Extend this partial E-function to an E-function $(\Omega''_1, f''_1), (\Omega''_2, f''_2), \ldots$ and let ξ' be the point in $\Omega(E, C)$ determined by this E-function. Then $\xi' \in U \subseteq W$, because $(\Omega''_j, f''_j) = (\Omega_j, f_j)$ for $1 \leq j \leq ir + t - 1$ and $ir + t - 1 \geq k$. On the other hand, if $\theta_g(\xi') = \xi'$, then it follows from Lemma 8.4 that

$$x = f''_{ir+t}(g_1 g^{-i}) = f''_t(g_1) = f_t(g_1)$$

which is a contradiction. Therefore $\theta_g(\xi') \neq \xi'$.

Now assume that $x_1 = y_1$. Then, since we are assuming that $x_1 \neq z_r$, we have that $y_1 \neq z_r$ and consequently

$$\eta\gamma^i = y_{2t-1}y^*_{2t-2}\cdots y^*_2 y_1 z^*_r x_r \cdots z^*_1 x_1 z^*_r x_r \cdots \quad \cdots z^*_1 x_1$$

is an admissible path. By Lemma 8.4, we have $f_{ir+t}(g_1 g^i) = f'_t(g_1) = f_t(g_1)$. So, proceeding as above, but now using the element $g_1 g^i$ instead of $g_1 g^{-i}$, we obtain an element ξ' in U such that $\theta_g(\xi') \neq \xi'$.

We now consider the case where $x_1 = z_r$. In this case we have, since $\theta_g(\xi) = \xi$,

$$z_1 = f_1(x_1) = f_1(z_r) = f'_1(z_r) = x_r.$$

Analogously, if in addition $x_2 = z_{r-1}$, then $z_2 = x_{r-1}$. Proceeding in this way, we see that there is an admissible path $\nu = z^*_i x_i \cdots z^*_1 x_1$ such that

$$\gamma = \nu^*\gamma'\nu$$

and $\gamma' = z^*_{r-i}x_{r-i}\cdots z^*_{i+1}x_{i+1}$ is a closed path in (E, C) such that $x_{i+1} \neq z_{r-i}$. Write $g' = z_{r-i}^{-1}x_{r-i}\cdots z_{i+1}^{-1}x_{i+1}$ and $h = z_i^{-1}x_i\cdots z_1^{-1}x_1$, and observe that $\theta_h(\xi)$ is a fixed point of $\theta_{g'}$.

Let W be an open neighborhood of ξ, which we may assume to be contained in the domain of θ_h. By the above, there is a point ξ' in the open neighborhood $\theta_h(W)$ of $\theta_h(\xi)$ such that $\theta_{g'}(\xi') \neq \xi'$. Then $\theta_{h^{-1}}(\xi') \in W$ and $\theta_g(\theta_{h^{-1}}(\xi')) \neq \theta_{h^{-1}}(\xi')$.

This completes the proof of the result in the case where g is a reduced word of the form $z_r^{-1}x_r z_{r-1}^{-1}x_{r-1}\cdots z_1^{-1}x_1$. Now assume that

$$g = x_r z_{r-1}^{-1}x_{r-1}\cdots z_1^{-1}x_1 z_0^{-1}$$

is a reduced word in $\mathbb{F}$, with $x_1, \ldots, x_r, z_0, \ldots, z_{r-1} \in E^1$ and $r \geq 1$. Assume that $\theta_g(\xi) = \xi$ for all $\xi \in V$, where V is a non-empty open subset of $\Omega(E, C)$ such that $V \subseteq \mathrm{Dom}(\theta_g)$. In particular this implies that $\mathrm{Dom}(\theta_g) \neq \emptyset$, so that $s(z_0) = s(x_1)$, $z^*_{r-1}x_{r-1}\cdots z^*_1 x_1$ is an admissible path, and $s(x_r) = s(z_{r-1})$. Since θ_g has fixed points we have $r(z_0) = r(x_r)$. By the same reason, either $x_r = z_0$ or x_r and z_0 belong to different elements of $C_{r(x_r)}$. Indeed, if $z_r \neq z_0$, and x_r and z_0 belong to $X \in C_{r(z_0)}$, then

$$V \subseteq \theta_{x_r}(\Omega(E,C)_{s(x_r)}) \cap \theta_{z_0}(\Omega(E,C)_{s(z_0)}) = H_{x_r} \cap H_{z_0} = \emptyset,$$

which is a contradiction.

Assume first that $x_r = z_0$. Then for $g' = z_{r-1}^* x_{r-1} \cdots z_1^* x_1$, we have that $g' \neq 1$ and that the set of fixed points of $\theta_{g'}$ contains the non-empty open set $\theta_{z_0^{-1}}(V)$. So we arrive to a contradiction by the first part of the proof.

If z_0 and x_r belong to different elements of $C_{r(z_0)}$, then set

$$g'' := z_0^{-1} x_r z_{r-1}^{-1} x_{r-1} \cdots z_1^{-1} x_1.$$

Then the set of fixed points of $\theta_{g''}$ contains the non-empty open set $\theta_{z_0^{-1}}(V)$, and we arrive again to a contradiction. This concludes the proof of the theorem. □

Remark 10.6. Let $w \in E^{0,1}$. We may define an entry based on w as a non-trivial admissible path of the form $x_{2t-1}x_{2t-2}^* \cdots x_4^* x_3 x_2^* x_1$ such that $s(x_1) = w$, and $|X| \geq 2$ for some $X \in C_{r(x_{2t-1})}$ such that $X \neq X_{x_{2t-1}}$. Every vertex $w \in E^{0,1}$ such that there is no entry based on w determines an isolated point in $\Omega(E,C)$. As another possible source of isolated points in $\Omega(E,C)$ we mention vertices $w \in E^{0,1}$ such that $|s^{-1}(w)| = 1$. For instance, in Example 9.6 there is a dense set of isolated points in the space X_v, namely all the points (in the representation provided by Figure 5) out of the union of the top row and the left-most column. Observe that however all three vertices in $E^{0,1}$ have entries based on them in that example. In many cases, $\Omega(E,C)$ is a Cantor set, that is, a metrizable, zero-dimensional compact set without isolated points.

We can now prove an analogue to the uniqueness theorem for graph C*-algebras with property (L). It applies to a *reduced* version of $\mathcal{O}(E,C)$, defined as follows:

Definition 10.7. Let (E,C) be a finite bipartite separated graph. Then we denote by $\mathcal{O}^r(E,C)$ the C*-algebra defined by the *reduced crossed product* of $C(\Omega(E,C))$ by $\mathbb{F}$:

$$\mathcal{O}^r(E,C) = (C(\Omega(E,C)) \rtimes_\alpha^r \mathbb{F}.$$

There exists a canonical *faithful* conditional expectation $E\colon \mathcal{O}^r(E,C) \to C(\Omega(E,C))$, see [22, Section 2].

Recall the following standard definition.

Definition 10.8. A C*-algebra satisfies property (SP) (for *small projections*) in case every nonzero hereditary C*-subalgebra contains a nonzero projection. Equivalently, for every nonzero positive element a in A there is $x \in A$ such that x^*ax is a nonzero projection.

Theorem 10.9. *Let (E,C) be a finite bipartite separated graph satisfying condition (L). Then the C*-algebra $\mathcal{O}^r(E,C)$ satisfies property* (SP). *More precisely, given a nonzero positive element c in $\mathcal{O}^r(E,C)$, there is an element $x \in \mathcal{O}^r(E,C)$ such that x^*cx is a nonzero projection in $C(\Omega(E,C))$. In particular every nonzero ideal of $\mathcal{O}^r(E,C)$ contains a nonzero projection of $C(\Omega(E,C))$.*

Proof. Since $\Omega(E,C)$ is topologically free when (E,C) satisfies condition (L) (Theorem 10.5), the same proof used in [5, Theorem 4.9] (which is based on [26, Proposition 2.4]) applies here. □

We can also derive a purely algebraic consequence. Note that the consideration of crossed products enable us to give a basically unified treatment of the purely algebraic case and the C*-case.

Recall that for a given field with involution K and a compact zero-dimensional topological space X, we denote by $C_K(X)$ the *-algebra of continuous functions $f\colon X\to K$, where K is given the discrete topology. Note that $C_{\mathbb{C}}(X)$ is a dense *-subalgebra of $C(X)$.

Theorem 10.10. *Let (E,C) be a finite bipartite separated graph satisfying condition (L). Then the *-algebra $L^{\mathrm{ab}}(E,C)=C_K(\Omega(E,C))\rtimes_\alpha \mathbb{F}$ satisfies property* (SP). *More precisely, given a nonzero element a in $L^{\mathrm{ab}}(E,C)$, there is a projection h in $C_K(\Omega(E,C))$, a nonzero element $\lambda\in K$ and an element $g\in\mathbb{F}$ such that*

$$ha(e(g)\delta_g)h=\lambda h\delta_e.$$

In particular every nonzero ideal of $L^{\mathrm{ab}}(E,C)$ contains a nonzero projection of $C_K(\Omega(E,C))$.

Proof. If is sufficient to show the result for any crossed product $\mathcal{A}=C_K(X)\rtimes_{\theta^*}G$, where G is a discrete group, X is a zero-dimensional compact space, and θ is a topologically free partial action of G on X such that U_t is a clopen subset of X for all $t\in G$. For $g\in G$, we denote by $e(g)$ the characteristic function of U_g, so that the ideals D_g associated to the induced action θ^* of G on $C_K(X)$ are given by $D_g=e(g)C_K(X)$. Recall that, setting $\alpha=\theta^*$, we have $\alpha_g(e(g^{-1})f)(\xi)=(e(g^{-1})f)(\theta_{g^{-1}}(\xi))=f(\theta_{g^{-1}}(\xi))$ for $\xi\in U_t$ and $f\in C_K(X)$.

We will show the statement for $\mathcal{A}=C_K(X)\rtimes_{\theta^*}G$ as above by using a slight modification of the arguments presented in [26, Proposition 2.4]. Let a be a nonzero element of $\mathcal{A}$. Multiplying a on the right by a suitable element of the form $e(g)\delta_g$, for $g\in G$, we obtain an element $c:=a(e(g)\delta_g)$ with $c_e\neq 0$, where $c=\sum_{t\in T}c_t\delta_t$ for a finite subset T of G. Note that $c_e=\sum_{i=1}^n\lambda_i\chi_{W_i}$, for some non-zero scalars $\lambda_i\in K$ and pairwise disjoint non-empty clopen subsets W_i of X. Setting $\lambda:=\lambda_1$, and using the fact that X is zero-dimensional, we obtain as in the proof of [26, Proposition 2.4] a clopen subset V contained in W_1 such that, putting $h:=\chi_V$, we have

$$h(c_t\delta_t)h=0,\qquad \text{for all } t\in T\setminus\{e\}.$$

Since $V\subseteq W_1$ we thus get $hch=hc_eh=\lambda h\delta_e$, so that we obtain

$$ha(e(g)\delta_g)h=hch=\lambda h\delta_e\,,$$

as desired. □

Remark 10.11. Observe that, in the conditions of Theorem 10.10, we also have that every nonzero one-sided ideal of $L^{\mathrm{ab}}(E,C)$ contains a nonzero idempotent. Indeed if a is a nonzero element of $L^{\mathrm{ab}}(E,C)$, and $ha(e(g)\delta_g)h=\lambda h\delta_e$ for some nonzero projection h in $C_K(X)$, some $g\in G$ and some $\lambda\in K\setminus\{0\}$, then $a[\lambda^{-1}(e(g)\delta_g)h]$ is a nonzero idempotent belonging to the right ideal generated by a, and analogously $[\lambda^{-1}(e(g)\delta_g)h]a$ is a nonzero idempotent in the left ideal generated by a.

Acknowledgements

It is a pleasure to thank Ken Goodearl and Takeshi Katsura for their useful comments.

References

[1] F. Abadie, *Enveloping actions and Takai duality for partial actions*, J. Funct. Anal. **197** (2003), 14–67.

[2] G. Abrams, G. Aranda Pino, *The Leavitt path algebra of a graph*, J. Algebra **293** (2005), 319–334.

[3] P. Ara, The realization problem for von Neumann regular rings, *Ring Theory 2007. Proceedings of the Fifth China-Japan-Korea Conference*, (eds. H. Marubayashi, K. Masaike, K. Oshiro, M. Sato); World Scientific, 2009, pp. 21–37.

[4] P. Ara, *Purely infinite simple reduced C*-algebras of one-relator separated graphs*, J. Math. Anal. Appl. **393** (2012), 493–508.

[5] P. Ara, R. Exel, T. Katsura, *Dynamical systems of type (m, n) and their C*-algebras*, Ergodic Theory and Dynamical Systems, to appear.

[6] P. Ara, A. Facchini, *Direct sum decompositions of modules, almost trace ideals, and pullbacks of monoids*, Forum Math. **18** (2006), 365–389.

[7] P. Ara, K. R. Goodearl, *C*-algebras of separated graphs*, J. Funct. Anal. **261** (2011), 2540–2568.

[8] P. Ara, K. R. Goodearl, *Leavitt path algebras of separated graphs.* J. reine angew. Math. **669** (2012), 165–224.

[9] P. Ara, K. R. Goodearl, *A wild refinement monoid*, in preparation.

[10] P. Ara, M. A. Moreno, E. Pardo, *Nonstable K-theory for graph algebras*, Algebr. Represent. Theory **10** (2007), 157–178.

[11] T. Bates, J. Hong, I. Raeburn and W. Szymański, *The ideal structure of the C*-algebras of infinite graphs*, Illinois J. Math. **46** (2002), 1159–1176.

[12] T. Bates, D. Pask, I. Raeburn and W. Szymański, *The C*-algebras of row-finite graphs*, New York J. Math. **6** (2000), 307–324 (electronic).

[13] B. BLACKADAR, "K-Theory for Operator Algebras", Second Edition, M.S.R.I. Publications, vol. 5, Cambridge Univ. Press, Cambridge, 1998.

[14] B Brenken, Z. Niu, *The C*-algebra of a partial isometry*, Proc. Amer. Math. Soc. **140** (2012), 199–206.

[15] L. G. Brown, *Ext of certain free product C^*-algebras*, J. Operator Theory **6** (1981), 135–141.

[16] G. M. Bergman, *Modules over coproducts of rings*, Trans. Amer. Math. Soc. **200** (1974) 1–32.

[17] G. M. Bergman, *Coproducts and some universal ring constructions*, Trans. Amer. Math. Soc. **200** (1974), 33–88.

[18] J. Cuntz and W. Krieger, *A class of C^*-algebras and topological Markov chains*, Inventiones Math. **56** (1980), 251–268.

[19] W. Dicks, T. Schick, *The spectral measure of certain elements of the complex group ring of a wreath product*, Geom. Dedicata **93** (2002), 121–137.

[20] M. Dokuchaev, R. Exel, *Associativity of crossed products by partial actions, enveloping actions and partial representations*, Trans. Amer. Math. Soc. **357** (2005), 1931–1952.

[21] M. Enomoto, Y. Watatani, *A graph theory for C*-algebras*, Math. Japon. **25** (1980), 435–442.

[22] R. Exel, *Amenability for Fell bundles*, J. Reine Angew. Math. **492** (1997), 41–73, [arXiv:funct-an/9604009].

[23] R. Exel, *Partial representations and amenable Fell bundles over free groups*, Pacific J. Math. **192** (2000), 39–63.

[24] Ruy Exel, *Inverse semigroups and combinatorial C*-algebras*, Bull. Braz. Math. Soc. (N.S.) **39** (2008), 191–313.

[25] R. Exel and M. Laca, *Cuntz–Krieger algebras for infinite matrices*, J. reine angew. Math. **512** (1999), 119–172, [arXiv:funct-an/9712008].

[26] R. Exel, M. Laca, J. Quigg, *Partial dynamical systems and C*-algebras generated by partial isometries*, J. Operator Theory 47 (2002), 169–186.

[27] C. Farthing, P. Muhly, and T. Yeend, *Higher-rank graph C*-algebras: an inverse semigroup and groupoid approach*, Semigroup Forum **71** (2005), 159–187.

[28] N. Fowler, M. Laca, and I. Raeburn, *The C*-algebras of infinite graphs*, Proc. Amer. Math. Soc. **128** (2000), 2319–2327.

[29] D. Gonçalves, D. Royer, Leavitt path algebras as partial skew group rings, arXiv:1202.2704v1 [math.RA].

[30] R. I. Grigorchuk, P. Linnell, T. Schick, A. Zuk, *On a question of Atiyah*, C. R. Acad. Sci. Paris Sér. I Math. **331** (2000), 663–668.

[31] P. Freyd, *Redei's finiteness theorem for commutative semigroups*, Proc. Amer. Math. Soc. **19** (1968), 1003.

[32] K. R. Goodearl, *Leavitt path algebras and direct limits*, Rings, modules and representations, 165–187, Contemp. Math., **480**, Amer. Math. Soc., Providence, RI, 2009.

[33] R. Hancock, I. Raeburn, *The C*-algebras of some inverse semigroups*, Bull. Austral. Math. Soc. **42** (1990), 335–348.

[34] D. Kerr, *C*-algebras and topological dynamics: finite approximation and paradoxicality*, CRM Advanced course books, Birkhauser, to appear; http://www.math.tamu.edu/∼kerr/.

[35] D. Kerr, P. W. Nowak, *Residually finite actions and crossed products*, Ergodic Theory and Dynamical Systems **32** (2012), 1585–1614.

[36] A. Kumjian and D. Pask, *C^*-algebras of directed graphs and group actions*, Ergodic Theory Dynam. Systems **19** (1999), 1503–1519.

[37] A. Kumjian, D. Pask, I. Raeburn, *Cuntz-Krieger algebras of directed graphs*, Pacific J. Math. **184** (1998), 161–174.

[38] A. Kumjian, D. Pask, I. Raeburn and J. Renault, *Graphs, groupoids, and Cuntz-Krieger algebras*, J. Funct. Anal. **144** (1997), 505–541.

[39] W. G. Leavitt. *The module type of a ring.* Trans. Amer. Math. Soc. **103** (1962) 113–130.

[40] K. McClanahan, *C^*-algebras generated by elements of a unitary matrix*, J. Funct. Anal. **107** (1992), 439–457.

[41] K. McClanahan, *K-theory and* Ext*-theory for rectangular unitary C^*-algebras*, Rocky Mountain J. Math. **23** (1993), 1063–1080.

[42] K. McClanahan, *Simplicity of reduced amalgamated products of C*-algebras*, Canad. J. Math. **46** (1994), 793–807.

[43] , ———, *K-theory for partial crossed products by discrete groups*, J. Funct. Anal. **130** (1995), 77–117.

[44] A. L. T. Paterson, *Graph inverse semigroups, groupoids and their C*-algebras*, J. Operator Theory **48** (2002), 645–662.

[45] I. Raeburn, Graph algebras. CBMS Regional Conference Series in Mathematics, 103. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2005.

[46] I. Raeburn, A. Sims and T. Yeend, *The C*-algebras of finitely aligned higher-rank graphs*, J. Funct. Anal. **213** (2004), 206–240.

[47] I. Raeburn, M. Tomforde and D. P. Williams, *Classification theorems for the C*-algebras of graphs with sinks*, Bull. Austral. Math. Soc. **70** (2004), 143–161.

[48] I. Raeburn and W. Szymański, *Cuntz-Krieger algebras of infinite graphs and matrices*, Trans. Amer. Math. Soc. **356** (2004), 39–59 (electronic).

[49] M. Rørdam, F. Larsen, N.J. Laustsen, "An Introduction to K-Theory for C^*-Algebras", Cambridge University Press, LMS Student Texts 49, 2000.

[50] M. Rørdam, A. Sierakowski, *Purely infinite C*-algebras arising from crossed products*, Ergodic Theory and Dynamical Systems **32** (2012), 273–293.

[51] A. Tarski, Cardinal Algebras. With an Appendix: Cardinal Products of Isomorphism Types, by Bjarni Jónsson and Alfred Tarski. Oxford University Press, New York, N. Y., 1949.

[52] M. Tomforde, *A unified approach to Exel-Laca algebras and C*-algebras associated to graphs*, J. Operator Theory **50** (2003), 345–368.

[53] J. K. Truss, *The failure of cancellation laws for equidecomposability types*, Canad. J. Math. **42** (1990), 590–606.

[54] S. Wagon, The Banach-Tarski paradox. With a foreword by Jan Mycielski. Corrected reprint of the 1985 original. Cambridge University Press, Cambridge, 1993.

[55] F. Wehrung, *Embedding simple commutative monoids into simple refinement monoids*, Semigroup Forum **56** (1998) 104–129.

Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Bellaterra (Barcelona), Spain.
E-mail address: `para@mat.uab.cat`

Departamento de Matemática, Universidade Federal de Santa Catarina, 88010-970 Florianópolis SC, Brazil.
E-mail address: `exel@mtm.ufsc.br`